





Bk

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 213/00		A2	(11) International Publication Number: WO 00/29378
			(43) International Publication Date: 25 May 2000 (25.05.00)
(21) International Application Number: PCT/EP99/08765		(74) Agent: BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent & Trademark Department, CH-4002 Basel (CH).	
(22) International Filing Date: 15 November 1999 (15.11.99)			
(30) Priority Data: 2298/98 17 November 1998 (17.11.98) CH 797/99 29 April 1999 (29.04.99) CH		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(71) Applicant (for all designated States except AT US): NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, D-4058 Basel (CH).			
(71) Applicant (for AT only): NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT M.B.H. [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT).			
(72) Inventors; and (75) Inventors/Applicants (for US only): GOEBEL, Thomas [DE/DE]; Schönaustrasse 1, D-79540 Lörrach (DE). HUMBERT-DROZ, Eliane [CH/CH]; Route de Noreaz 108, CH-1772 Ponthaux (CH). SCHWARZENBACH, Maurizio [CH/CH]; Unterer Brunnacher 22, CH-4423 Ramllinsburg (CH).		Published Without international search report and to be republished upon receipt of that report.	

(54) Title: PESTICIDES



(57) Abstract

Novel pesticides of formula (I), wherein the substituents R, R₁, R₂, R₂', T, U, X and Y are as defined in claim 1, are described. Also described are compositions suitable for use as parasitocides comprising those compounds as active ingredient and to methods of controlling parasites that are based on the administration of those compounds or compositions, and to the use of the said compounds and compositions in a method of controlling parasites and in the manufacture of pesticides for use against parasites. Also described are intermediates of formula (XX) wherein R₁, R₂, R₂', T, U, X and Y are as defined in claim 1 and Hal is halogen. The latter also exhibit parasitocidal activity and are suitable for the preparation of the compounds of formula (I).

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						


Pesticides

The present invention relates to novel pesticides of the formula (I) below having improved action against parasites; to compositions suitable for use as parasiticides comprising those compounds as active ingredient and to methods of controlling parasites that are based on the administration of those compounds or compositions, and to the use of the said compounds and compositions in a method of controlling parasites and in the manufacture of pesticides for use against parasites. Also described are intermediates of formula (XX) which themselves have parasitocidal activity and are excellently suitable for the preparation of compounds of formula (I).

Numerous pesticides are known that can be used in controlling parasites on warm-blooded animals. The control is effected principally by two different methods; either by way of contact action by topical and therefore external treatment of the host animal or systemically, that is to say by oral, transdermal or percutaneous administration to the host animal and ingestion of the active ingredient by the parasites *via* the blood of the host animal.


Far fewer substances are available for systemic use than for topical application, because only substances that have a systemic action and are well tolerated by the host animal can be used.

Compounds having as characteristic structural element the sub-structure of formula (II)



wherein R_1 , X, Y, T and U are as defined for formula (I) hereinbelow, form a very interesting class of substances on account of their pronounced topical and systemic action.

A prominent individual example is nitenpyram, a compound of formula (III)



Nitenpyram and other examples of this class of substances are disclosed, together with their preparation, in EP 0 302 389. Those compounds are described as pesticides having very pronounced insecticidal activity. Further examples of this class of substances are, for example, the subjects of the following patent applications: European published specifications Nos. 285 985; 302 833; 376 279; 471 372; 364 844; 493 369; 381 130; 529 680; 163 855; 375 907; 259 738; 386 565; 383 091 and 590 425; US Patents 5 063 236; 5 302 605 and 4 742 060; and also DE-4 207 604; GB-2 228 003 and WO 93/24002.

Nitenpyram and other examples of this class of substances that have the said structural element of formula (II) are extremely effective when administered as contact pesticides, for example externally, that is to say topically, to an infested host animal where they come into direct contact with the parasites. They also, however, exhibit a good systemic immediate action when they are administered to the infested host animal orally, parenterally, via injection or *via* implant.

The action, which is pronounced *per se*, has a serious disadvantage, however, in that it has been found that while the compounds have a high initial action, their action falls off rapidly only a short time after administration. This can be observed particularly clearly after systemic administration and can be monitored by reference to the bioavailability. Blood level measurements show that in many cases high blood levels are achieved even after a few minutes or, more rarely, after a few hours, but these levels then fall within a few hours, at best within a few days, and therefore fall to below an effective concentration much too rapidly.

In order to eliminate this shortcoming, numerous, but unfortunately unsuccessful, experiments have already been carried out. For example, it has been shown that a prolongation of the systemic action by increasing the dose can be achieved only to a limited extent. If, for example, depots sufficiently large for the active ingredient to be released over several weeks were to be placed under the skin or in the muscles, then the amounts to be injected


or implanted would have to be so large that they would no longer be tolerated by the host animal; local irritation, skin eruptions and painful areas develop. This solution, possible *per se*, therefore fails on practical and, of course, also ethical grounds. Similarly, it has been found that a long-term action is not achievable by an increased oral dose.

When the known compounds having the structural element of formula (II) are administered, it is principally observed that a major part of the substance exhibits its full action only over a short period of time immediately after administration and thereafter the action very rapidly declines. This has serious consequences for preparations for use in veterinary medicine, for example for tablets, injections or for treatment using the pour-on or spot-on method. Because of the short duration of action it is necessary to repeat treatments at short intervals, which means that the keeper of the animal must either repeat the treatment himself, or have it carried out by a veterinary surgeon, at short intervals. Such an intensive treatment programme requires a high degree of discipline, however, and, as experience has shown, after only a short time gives rise to stress on the part of the animal and on the part of the animal's keeper, which not infrequently results in aversion to the treatment and leads to its premature discontinuation.

Prolongation of the action of this inherently extremely effective class of substances has therefore long been a desirable but apparently unattainable goal. The problem underlying the present invention was to achieve that goal and provide substances suitable for use as pesticides having significantly improved properties, especially having a pronounced long-term action.

By the provision of the compounds of formula (I) below it has now, surprisingly, been possible for compounds having the structural element of formula (II) to be modified chemically in such a manner that a high degree of long-term bioavailability after administration is achieved without it being necessary to accept adverse effects as a result.

The new, improved compounds are compounds of formula (I) below




wherein

- R₁ is hydrogen or a radical from the group C₁-C₄alkyl, formyl, C₁-C₆alkylcarbonyl, C₁-C₄-alkylsulfonyl, aryl, arylsulfonyl, arylcarbonyl, heterocyclyl and heterocyclyl-substituted C₁-C₆alkyl, which radical is unsubstituted or mono- or poly-substituted by identical or different substituents; the said substituents being C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄alkylthio, C₁-C₄haloalkyl, halogen, hydroxy, cyano, nitro, amino, C₁-C₄alkylamino, (C₁-C₄-alkyl)₂amino, alkoxycarbonyl, C₁-C₄alkylsulfonyl and arylsulfonyl;
- X is CH or N;
- Y is an electron-withdrawing radical, preferably cyano, nitro or C₁-C₆haloalkyl-carbonyl, especially CO-CF₃;
- T has the meanings of R₁ or together with U forms a C₁-C₄alkylene bridge which is unsubstituted or substituted by a radical R₁, or T and U together with the group -N-C-N- form a saturated or unsaturated 5- or 6-membered heterocyclic ring which may in addition contain as further hetero atom O or S or the hetero group -N(C₁-C₆alkyl)-;
- U is hydrogen or C₁-C₆alkyl, preferably hydrogen, methyl or ethyl;
- R₂ is hydrogen or C₁-C₆alkyl; R₂' is hydrogen or C₁-C₆alkyl; and
- R is C₁-C₂₀alkyl, C₂-C₂₀alkenyl, C₂-C₆alkynyl or heterocyclyl, each of those radicals being unsubstituted or substituted by one or more identical or different substituents, the said substituents being selected from the group halogen, cyano, nitro, hydroxy, C₁-C₆alkoxy, C₁-C₆alkylthio, C₁-C₆haloalkyl, C₁-C₆haloalkoxy and phenyl; or is C₃-C₇cycloalkyl that is unsubstituted or mono- or poly-substituted by identical or different substituents selected from halogen, cyano, nitro, hydroxy, C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, C₁-C₆alkoxy, C₁-C₆alkylthio, C₁-C₆haloalkyl, C₁-C₂₀haloalkoxy and phenyl; wherein each phenyl moiety is itself unsubstituted or mono- or poly-substituted by identical or different substituents selected from halogen, cyano, nitro, hydroxy, C₁-C₆alkyl, C₁-C₆alkoxy, C₁-C₆alkylthio, C₁-C₆haloalkyl and C₁-C₂₀haloalkoxy; or is phenyl phenoxyphenyl each of which is unsubstituted or mono- or poly-substituted by identical or different substituents selected from halogen, cyano, nitro, hydroxy, C₁-C₆alkyl, C₁-C₆alkoxy, C₁-C₆alkylthio, C₁-C₆haloalkyl, and C₁-C₂₀haloalkoxy.

Within the scope of formula (I) above, preference is given to compounds wherein R is C₁-C₂₀alkyl, C₂-C₂₀alkenyl or C₂-C₆alkynyl, each of those radicals being unsubstituted or mono- or poly-substituted by identical or different substituents, the said substituents being selected from the group halogen, cyano, nitro, hydroxy, C₁-C₆alkoxy, C₁-C₆alkylthio, C₁-C₆haloalkyl, C₁-C₆haloalkoxy and phenyl; or is C₃-C₇cycloalkyl that is unsubstituted or mono- or poly-substituted by identical or different substituents selected from halogen, cyano, nitro, hydroxy, C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, C₁-C₆alkoxy, C₁-C₆alkylthio, C₁-C₆haloalkyl, C₁-C₂₀haloalkoxy and phenyl; or is phenyl phenoxyphenyl each of which is unsubstituted or mono- or poly-substituted by identical or different substituents selected from halogen, cyano, nitro, hydroxy, C₁-C₆alkyl, C₁-C₆alkoxy, C₁-C₆alkylthio, C₁-C₆haloalkyl, C₁-C₂₀haloalkoxy.

Within the group of compounds of formula (I) wherein R is C₃-C₇cycloalkyl that is mono- or poly-substituted by identical or different substituents selected from halogen, cyano, nitro, hydroxy, C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, C₁-C₆alkoxy, C₁-C₆alkylthio, C₁-C₆haloalkyl, C₁-C₂₀haloalkoxy and phenyl; wherein the phenyl moiety is itself unsubstituted or mono- or poly-substituted by identical or different substituents selected from halogen, cyano, nitro, hydroxy, C₁-C₆alkyl, C₁-C₆alkoxy, C₁-C₆alkylthio, C₁-C₆haloalkyl and C₁-C₂₀haloalkoxy, special preference on account of their pronounced activity is given to those compounds in which the C₃-C₇cycloalkyl radical is substituted by one substituent and in the 1-position.

By the introduction of the side chain of formula (IV)



wherein R₂, R₂' and R are as defined for formula (I), it has now been possible to prepare substances that exhibit simultaneously a number of long sought and very desirable properties and are now actually suitable for practical use:

1. The compounds of formula (I) have a high level of activity against arthropods, especially blood-sucking insects.
2. They exhibit excellent tolerability when administered systemically and topically to a host animal.

3. They are distinguished by an appreciably longer duration of action in comparison with known compounds having the structural element of formula (II), which can readily be demonstrated by reference to the mortality of the parasites on the host animal.
4. They can be handled satisfactorily from the standpoint of formulation technology and have adequate storage stability.

It was not to be predicted that the chemical modification carried out here would result in these advantageous properties and would be able to impart to the novel compounds of formula (I) these positive long-term properties.

In the context of the present invention, the definitions of the substituents are to be understood as follows: each of the substituents indicated under formula (I) that can itself be poly-substituted is substituted by either identical or different substituents, that is to say multiple substitutions are to be interpreted as meaning that identical or different substituents can be present simultaneously on the same radical. For example, a radical poly-substituted by halogen may have either several identical halogen atoms or several different halogen atoms. Multiple substitutions are to be interpreted accordingly for other radicals.

The alkyl groups appearing in the definitions of substituents in terms such as alkyl, alkyl-carbonyl, alkylsulfonyl, alkoxy, alkylthio, haloalkyl, alkylamino, dialkylamino, haloalkyl-carbonyl, etc. may, according to the number of carbon atoms, be straight-chain or branched and are, for example, methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, dodecyl, tetradecyl, hexadecyl, octadecyl or eicosyl, and the branched isomers thereof, e.g. isopropyl, isobutyl, sec-butyl, tert-butyl, isopentyl, neopentyl or isohexyl. Alkoxy, haloalkyl, haloalkylcarbonyl and haloalkoxy radicals are derived from the mentioned alkyl groups and accordingly are partially or fully halogenated radicals; poly-halogenated radicals carry identical or different halogen atoms.

The terms halo and halogen denote halogen atoms and generally denote fluorine, chlorine, bromine or iodine, here preferably fluorine or chlorine, as substituent of an alkyl group especially fluorine and as substituent of a heterocycle especially chlorine.

Examples of haloalkyl – as a group *per se* and as a structural element of other groups and compounds, such as haloalkoxy – are methyl mono- to tri-substituted by fluorine, chlorine and/or bromine, such as CHF_2 or CF_3 ; ethyl mono- to penta-substituted by fluorine, chlorine

and/or bromine, such as CH_2CF_3 , CF_2CF_3 , CF_2CCl_3 , CF_2CHCl_2 , CF_2CHF_2 , CF_2CFCl_2 , CF_2CHBr_2 , CF_2CHClF , CF_2CHBrF or CClFCHClF ; propyl or isopropyl mono- to hepta-substituted by fluorine, chlorine and/or bromine, such as $\text{CH}_2\text{CHBrCH}_2\text{Br}$, $\text{CF}_2\text{CHFCH}_2\text{F}$, $\text{CH}_2\text{CF}_2\text{CF}_3$ or $\text{CH}(\text{CF}_3)_2$; and butyl mono- to nona-substituted by fluorine, chlorine and/or bromine, or one of its isomers, such as $\text{CF}(\text{CF}_3)\text{CHFCF}_3$ or $\text{CH}_2(\text{CF}_2)_2\text{CF}_3$.

In the context of the present invention, Het and heterocyclyl are to be understood as meaning aliphatic or aromatic cyclic radicals that contain at least one oxygen, sulfur or nitrogen atom. Five- and six-membered heterocycles are preferred. Heterocyclyl accordingly typically includes substituents such as dioxolanyl, pyrrolidinyl, piperidinyl, morpholinyl, pyridyl, pyrrolyl, furyl, thienyl, imidazolyl, tetrahydrofuryl, tetrahydropyranyl, dihydrofuryl, dihydropyranyl, isoxazolyl, oxazolyl, thiazolyl, oxazolynyl, oxazolidinyl, imidazolynyl, imidazolidinyl and dioxanyl. Special preference is given to those which are unsubstituted or contain one or two halogen atoms, halogen here being fluorine, chlorine or bromine, but especially chlorine. Of those heterocyclic radicals special mention should be made of pyridyl, thiazolyl and tetrahydrofuryl. More especially preferred sub-groups of formula (1) contain as heterocyclyl radicals 5,6-dichloro-pyridin-3-yl, 6-chloro-pyridin-3-yl, 2-chlorothiazol-5-yl and tetrahydrofuran-3-yl.

Aryl by itself or as part of a substituent, for example arylsulfonyl, arylcarbonyl or aralkyl, is phenyl or naphthyl, preferably phenyl.

Alkenyl is, in each case giving due consideration to the number of carbon atoms contained in the group in question, either straight-chain, e.g. vinyl, 1-methylvinyl, allyl, 1-butenyl or 2-hexenyl, or branched, e.g. isopropenyl. Alkynyl is, in each case giving due consideration to the number of carbon atoms contained in the group in question, either straight-chain, e.g. propargyl, 2-butenyl or 5-hexynyl, or branched, e.g. 2-ethynylpropyl or 2-propargylisopropyl. $\text{C}_3\text{-C}_7$ Cycloalkyl is cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl.

Typical $\text{C}_1\text{-C}_4$ alkylene bridges are $-\text{CH}_2-$, $-\text{CH}_2\text{-CH}_2-$, $-\text{CH}_2\text{-CH}_2\text{-CH}_2-$, $-\text{CH}_2\text{-CH}_2\text{-CH}_2\text{-CH}_2-$, $-\text{CH}(\text{CH}_3)\text{-CH}_2-$, $-\text{CH}(\text{C}_2\text{H}_5)\text{-CH}_2-$, $-\text{CH}_2\text{-CH}(\text{CH}_3)\text{-CH}_2-$, $-\text{CH}(\text{CH}_3)\text{-CH}(\text{CH}_3)-$, $-\text{C}(\text{CH}_3)_2-$, $-\text{C}(\text{CH}_3)(\text{C}_2\text{H}_5)-$ and $-\text{C}(\text{C}_2\text{H}_5)_2-$.

Compounds of formula (I) having at least one basic centre may form acid addition salts with strong acids. Physiologically tolerable acid addition salts are of special interest.

An interesting sub-group within the compounds according to the invention is formed by compounds of formula (I) wherein

Y is NO₂;

R₁ is hydrogen or a radical from the group C₁-C₄alkyl, formyl, C₁-C₆alkylcarbonyl, C₁-C₄-alkylsulfonyl, aryl, arylsulfonyl, arylcarbonyl, heterocyclyl and heterocyclyl-substituted C₁-C₆alkyl, which radical is unsubstituted or mono- or poly-substituted by identical or different substituents; the said substituents being C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄alkylthio, C₁-C₄haloalkyl, halogen, hydroxy, cyano, nitro, amino, C₁-C₄alkylamino, (C₁-C₄-alkyl)₂amino, alkoxycarbonyl, C₁-C₄alkylsulfonyl and arylsulfonyl;

T has the meanings of R₁ or together with U forms a C₁-C₄alkylene bridge which is unsubstituted or substituted by a radical R₁; or T and U together with the group -N-C-N- form a saturated or unsaturated 5- or 6-membered heterocyclic ring which may in addition contain as further hetero atom O or S or the hetero group -N(C₁-C₆alkyl)-;

U is hydrogen or C₁-C₆alkyl, preferably hydrogen, methyl or ethyl; and R₂, R₂' and R are as defined for formula (I).

Another interesting group is formed by compounds of formula (I) wherein

R₁ is -CH₂-Het; X is CH; Y is NO₂;


Het is heterocyclyl that is unsubstituted or mono- or poly-substituted by identical or different substituents; the substituents being C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄alkylthio, C₁-C₄haloalkyl, halogen, hydroxy, cyano, nitro, amino, C₁-C₄alkylamino, (C₁-C₄alkyl)₂amino, alkoxycarbonyl, C₁-C₄alkylsulfonyl and arylsulfonyl;

T (1) is a radical from the group formyl, C₁-C₆alkylcarbonyl, C₁-C₄alkylsulfonyl, aryl, arylsulfonyl, arylcarbonyl, heterocyclyl and heterocyclyl-substituted C₁-C₆alkyl, which radical is unsubstituted or mono- or poly-substituted by identical or different substituents; the said substituents being C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkyl, halogen, hydroxy, cyano, nitro, amino, C₁-C₄alkylamino, (C₁-C₄alkyl)₂amino, C₁-C₄alkylsulfonyl and arylsulfonyl; or
 (2) T together with U forms a C₁-C₄alkylene bridge which is unsubstituted or substituted by a radical selected from the group C₁-C₄alkyl, formyl, C₁-C₆alkylcarbonyl, C₁-C₄alkylsulfonyl, aryl, arylsulfonyl, arylcarbonyl, heterocyclyl and heterocyclyl-substituted C₁-C₆-

alkyl; each radical from the said group itself being unsubstituted or substituted by C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkyl, halogen, hydroxy, cyano, nitro, amino, C₁-C₄-alkylamino, (C₁-C₄alkyl)₂amino, C₁-C₄alkylsulfonyl or arylsulfonyl; or
 (3) T and U together with the group -N-C-N- form a saturated or unsaturated 5- or 6-membered heterocyclic ring which may in addition contain as further hetero atom O or S or the hetero group -N(C₁-C₆alkyl)-;

U is hydrogen or C₁-C₆alkyl, preferably hydrogen, methyl or ethyl; and R₂, R₂' and R are as defined for formula (I).

Very especially preferred within the scope of formula (I), however, are compounds of formula (X)



wherein

R₁ is -CH₂-Het;


R is C₁-C₂₀alkyl, C₂-C₂₀alkenyl or C₂-C₆alkynyl, each of those radicals being unsubstituted or mono- or poly-substituted by identical or different substituents, the said substituents being selected from the group halogen, cyano, nitro, hydroxy, C₁-C₆alkoxy, C₁-C₆alkylthio, C₁-C₆haloalkyl, C₁-C₆haloalkoxy and phenyl; or is C₃-C₇cycloalkyl that is unsubstituted or mono- or poly-substituted by identical or different substituents selected from halogen, cyano, nitro, hydroxy, C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, C₁-C₆alkoxy, C₁-C₆alkylthio, C₁-C₆haloalkyl, C₁-C₂₀haloalkoxy and phenyl; or is phenyl phenoxy-phenyl each of which is unsubstituted or mono- or poly-substituted by identical or different substituents selected from halogen, cyano, nitro, hydroxy, C₁-C₆alkyl, C₁-C₆alkoxy, C₁-C₆alkylthio, C₁-C₆haloalkyl, C₁-C₂₀haloalkoxy;

T and U are each independently of the other hydrogen or C₁-C₆alkyl, preferably hydrogen, methyl or ethyl;

R₂ is hydrogen or C₁-C₆alkyl; R₂' is hydrogen or C₁-C₆alkyl; and

Het is heterocyclyl that is unsubstituted or mono- or poly-substituted by identical or different halogen atoms.


A further especially preferred sub-group of compounds of formula (I) is formed, on account of their pronounced activity, by compounds of formula (XI)



wherein

- Hal is halogen, preferably fluorine, chlorine or bromine and especially chlorine; and especially occupies the 6-position in the pyridine;
- X is CH or N and especially N;
- Y is an electron-withdrawing radical, preferably cyano, nitro or C₁-C₆haloalkyl-carbonyl, especially CO-CF₃; more especially nitro;
- T together with U forms a C₁-C₄alkylene bridge, preferably an ethylene bridge, which is preferably unsubstituted or substituted by methyl or ethyl; and R₂, R₂' and R are as defined for formula (I).

Equally preferred on account of their pronounced activity is a sub-group of compounds of formula (I) having the following formula (XII)



wherein

- Hal is halogen, preferably fluorine, chlorine or bromine and especially chlorine; and especially occupies the 2-position in the thiazole;
- X is CH or N and especially N;
- Y is an electron-withdrawing radical, preferably cyano, nitro or C₁-C₆haloalkyl-carbonyl, especially CO-CF₃; more especially nitro;
- T together with U forms one of the groups -CH₂-CH₂-, -CH₂-CH₂-CH₂-, -CH₂-O-CH₂- and -CH₂-N(CH₃)-CH₂-, wherein all methylene groups are unsubstituted or one of said methylene groups is substituted by methyl or ethyl; and R₂, R₂' and R are as defined in claim 1 for formula (I).

A preferred sub-group of compounds within the scope of formula (X) is formed by those compounds wherein U is methyl or ethyl.

A further preferred group of compounds within the scope of formula (X) is formed by those compounds wherein T is methyl or ethyl.

A further preferred sub-group of compounds within the scope of formula (X) is formed by those compounds wherein R₂ and R₂' are hydrogen, methyl or ethyl.

Especially preferred among the compounds within the scope of formula (X) and within the scope of preferred sub-groups mentioned above are those compounds in which the radical Het is pyridyl, thiazolyl or tetrahydrofuryl that is unsubstituted or mono- or di-substituted by halogen; especially 5,6-dichloro-pyridin-3-yl, 6-chloro-pyridin-3-yl, 2-chlorothiazol-5-yl and tetrahydrofuran-3-yl.

Within the scope of mentioned sub-groups, preference is given to compounds of formula (I) wherein R is C₁-C₂₀alkyl, C₂-C₂₀alkenyl or C₂-C₆alkynyl and especially straight-chain or branched C₆-C₂₀alkyl.

Especially preferred on account of its biological activity is any compound selected from the group of compounds 1.001; 1.008; 1.011; 1.012; 1.013; 1.014; 1.015; 1.018; 1.019; 1.020; 1.021; 1.022; 1.054; 1.055; 1.056; 1.057; 1.058; 1.059; 1.060; 1.061; 1.062; 1.063; 1.064; 1.065; 1.066; 1.067; 1.068; 1.069; 1.070; 1.071; 1.072; 1.073; 1.074; 1.075; 1.076; 1.077; 1.078; 1.079; 1.080; 1.081; 1.082; 1.083; 1.084; 1.085; 1.086 and 1.087.

Parasites in the context of the present invention are parasitic arthropods and, of those, especially blood-sucking insects. Insects of the following orders are included: *Lepidoptera*, *Coleoptera*, *Homoptera*, *Heteroptera*, *Diptera*, *Thysanoptera*, *Orthoptera*, *Anoplura*, *Siphonaptera*, *Mallophaga*, *Thysanura*, *Isoptera*, *Psocoptera* and *Hymenoptera*. Special mention should also be made, however, of pests that trouble human beings or animals and transmit pathogens, for example flies, such as *Musca domestica*, *Musca vetustissima*, *Musca autumnalis*, *Fannia canicularis*, *Sarcophaga carnaria*, *Lucilia sericata*, *Lucilia cuprina*, *Hypoderma bovis*, *Hypoderma lineatum*, *Chrysomya chloropyga*, *Dermatobia hominis*,

Cochliomyia hominivorax, *Gasterophilus intestinalis*, *Oestrus ovis*, *Callipora erythrocephala* (= blowfly), *Haematobia* (= hornfly) and mosquitos, and also blood-sucking pests, for example fleas, such as *Ctenocephalides felis* and *Ctenocephalides canis* (cat and dog fleas), *Xenopsylla cheopis*, *Pulex irritans*, *Dermatophilus penetrans*, lice, such as *Damalina ovis*, *Pediculus humanis*, stable flies and horseflies, such as *Stomoxys calcitrans*, *Haematopota pluvialis*, *Tabanus nigrovittatus*, *Chrysops caecutiens*, tabanids, tsetse flies, such as *Glossinia* species, and biting insects, more especially cockroaches, such as *Blatella germanic* or *Blatta orientalis*, *Periplaneta americana*. The said parasites attack warm-blooded animals, including farm animals, such as cows, pigs, sheep and goats, poultry, such as hens, turkeys and geese, animals bred for their fur, such as mink, foxes, chinchillas, rabbits and the like, as well as domestic animals and pets, such as cats and dogs, and even human beings do not escape attack.

Likewise, flea infestation in domestic animals and pets is a problem for the animal owner to which as yet only unsatisfactory solutions have been found. Owing to the complicated life cycle of the flea, none of the known methods of controlling fleas is totally satisfactory, especially since most of the known methods are aimed principally at controlling the fully grown fleas in the animal's coat and take no account at all of the various juvenile stages of the fleas, which live not only in the animal's coat but also on the floor, on carpets, on the animal's sleeping place, on chairs, in the garden and in all the other places with which the infested animal comes into contact. Flea treatment is generally expensive and must be continued for a prolonged period, success generally only being achieved when the treatment is applied not only to the infested animal, e.g. the dog or cat, but also simultaneously to any places frequented by the infested animal.

The compounds of formula (I) according to the invention can be used alone or in combination with other biocides. For example, in order to increase the effect they can be combined with pesticides having the same direction of action or in order to broaden the spectrum of action they can be combined with substances having a different direction of action. It may also be of advantage to add repelling substances, so-called repellents. Where it is desired to extend the spectrum of action to endoparasites, e.g. worms, the compounds of formula (I) are advantageously combined with substances having endoparasitocidal properties. They can, of course, also be used in combination with anti-bacterial agents. Since the

compounds of formula (I) are "adulticides", that is to say since they are effective especially against the fully grown stages of the target parasites, the addition of pesticides that are effective rather against the juvenile stages of the parasite may be very advantageous, since in that way the majority of parasites causing large-scale economic damage will be covered. Furthermore, a substantial contribution is made to avoiding the development of resistance. Some combinations may also lead to synergistic effects, that is to say the total amount of active ingredient applied can be reduced, which is desirable from the ecological standpoint. Preferred groups of combination partners and especially preferred combination partners are given below, it being possible for the combinations to comprise one or more of these partners in addition to a compound of formula (I).

Suitable mixing partners include biocides, for example the insecticides and acaricides listed below, which have various mechanisms of action and are well known to the person skilled in the art, for example chitin synthesis inhibitors, growth regulators; active ingredients that act in the same way as juvenile hormones; active ingredients that act as adulticides; broad spectrum insecticides; broad spectrum acaricides and nematocides; and also the well known anthelmintics and substances that repel insects and/or acarina, the afore-mentioned repellents or detachers.

Non-limiting examples of suitable insecticides and acaricides are:

(I) aldicarb;	(XV) deltamethrin;	(XXIX) mevinphos;
(II) azinphos-methyl;	(XVI) diflubenzuron;	(XXX) parathion;
(III) benfuracarb;	(XVII) endosulfan;	(XXXI) parathion-methyl;
(IV) bifenthrin;	(XVIII) ethiofencarb;	(XXXII) phosalone;
(V) buprofezin;	(XIX) fenitrothion;	(XXXIII) pirimicarb;
(VI) carbofuran;	(XX) fenobucarb;	(XXXIV) propoxur;
(VII) dibutylaminothio;	(XXI) fenvalerate;	(XXXV) teflubenzuron;
(VIII) cartap;	(XXII) formothion;	(XXXVI) terbufos;
(IX) chlorfluazuron;	(XXIII) methiocarb;	(XXXVII) triazamate;
(X) chlorpyrifos;	(XXIV) heptenophos;	(XXXVIII) abamectin;
(XI) cyfluthrin;	(XXV) imidacloprid;	(XXXIX) fenobucarb;
(XII) lambda-cyhalothrin;	(XXVI) isoprocarb;	(XL) tebufenozide;
(XIII) alpha-cypermethrin;	(XXVII) methamidophos;	(XLI) fipronil;
(XIV) zeta-cypermethrin;	(XXVIII) methomyl;	(XLII) beta-cyfluthrin;

- | | | |
|--|---------------------------|-------------------------|
| (XLIII) silafluofen; | (XLVI) fenazaquin; | (XLIX) nitenpyram; |
| (XLIV) fenpyroximate; | (XLVII) pyriproxyfen; | (L) NI-25, acetamiprid; |
| (XLV) pyridaben; | (XLVIII) pyrimidifen; | |
| (LI) avermectin B ₁ ; | | |
| (LII) an insect-active extract from a plant; | | |
| (LIII) a preparation containing insect-active nematodes; | | |
| (LIV) a preparation obtainable from <i>Bacillus subtilis</i> ; | | |
| (LV) a preparation containing insect-active fungi; | | |
| (LVI) a preparation containing insect-active viruses; | | |
| (LVII) AC 303 630; | (LXXX) chlorethoxyfos; | (CIV) fenamiphos; |
| (LVIII) acephat; | (LXXXI) chlormephos; | (CV) fenbutatinoxid; |
| (LIX) acrinathrin; | (LXXXII) cis-Res-methrin; | (CVI) fenothiocarb; |
| (LX) alanycarb; | (LXXXIII) clocythrin; | (CVII) fenpropathrin; |
| (LXI) alphamethrin; | (LXXXIV) clofentezin; | (CVIII) fenpyrad; |
| (LXII) amitraz; | (LXXXV) cyanophos; | (CIX) fenthion; |
| (LXIII) AZ 60541; | (LXXXVI) cycloprothrin; | (CX) fluazinam; |
| (LXIV) azinphos A; | (LXXXVII) cyhexatin; | (CXI) flucycloxuron; |
| (LXV) azinphos M; | (LXXXVIII) demeton M; | (CXII) flucythrinate; |
| (LXVI) azocyclotin; | (LXXXIX) demeton S; | (CXIII) flufenoxuron; |
| (LXVII) bendiocarb; | (XC) demeton-S-methyl; | (CXIV) flufenprox; |
| (LXVIII) bensultap; | (XCI) dichlofenthion; | (CXV) fonophos; |
| (LXIX) beta-cyfluthrin; | (XCII) dicliphos; | (CXVI) fosthiazat; |
| (LXX) BPMC; | (XCIII) diethion; | (CXVII) fubfenprox; |
| (LXXI) brofenprox; | (XCIV) dimethoat; | (CXVIII) HCH; |
| (LXXII) bromophos A; | (XCV) dimethylvinphos; | (CXIX) hexaflumuron; |
| (LXXIII) bufencarb; | (XCVI) dioxathion; | (CXX) hexythiazox; |
| (LXXIV) butocarboxime; | (XCVII) edifenphos; | (CXXI) iprobenfos; |
| (LXXV) butylpyridaben; | (XCVIII) emamectin; | (CXXII) isofenphos; |
| (LXXVI) cadusafos; | (XCIX) esfenvalerat; | (CXXIII) isoxathion; |
| (LXXVII) carbaryl; | (C) ethion; | (CXXIV) ivermectin; |
| (LXXVIII) carbophenothion; | (CI) ethofenprox; | (CXXV) lambda- |
| (LXXIX) chloethocarb; | (CII) ethoprophos; | cyhalothrin; |
| | (CIII) etrimphos; | (CXXVI) malathion; |

(CXXVII) mecarbam;	(CLIX) tebufenpyrad;	(CLXXXIX) fenoxycarb
(CXXVIII) mesulfenphos;	(CLX) tebupirimphos;	(CXC) chlorfenapyr or
(CXXIX) metaldehyd;	(CLXI) tefluthrin;	(CXCI) spinosad
(CXXX) metolcarb;	(CLXII) temephos;	
(CXXXI) milbemectin;	(CLXIII) terbam;	
(CXXXII) moxidectin;	(CLXIV) tetrachlor-	
(CXXXIII) naled;	vinphos;	
(CXXXIV) NC 184;	(CLXV) thiafenox;	
(CXXXV) omethoat;	(CLXVI) thiodicarb;	
(CXXXVI) oxamyl;	(CLXVII) thiofanox;	
(CXXXVII) oxydeme- thon	(CLXVIII) thionazin;	
M;	(CLXIX) thuringiensin;	
(CXXXVIII) oxydeprofos;	(CLXX) tralomethrin;	
(CXXXIX) permethrin;	(CLXXI) triarthen;	
(CXL) phenthoat;	(CLXXII) triazophos;	
(CXLI) phorat;	(CLXXIII) triazuron;	
(CXLII) phosmet;	(CLXXIV) trichlorfon;	
(CXLIII) phoxim;	(CLXXV) triflumuron;	
(CXLIV) pirimiphos M;	(CLXXVI) trimethacarb;	
(CXLV) pirimiphos A;	(CLXXVII) vamidothion;	
(CXLVI) promecarb;	(CLXXVIII) xylylcarb;	
(CXLVII) propaphos;	(CLXXIX) YI 5301/5302;	
(CXLVIII) prothiofos;	(CLXXX) zetamethrin;	
(CXLIX) prothoat;	(CLXXXI) DPX-MP062;	
(CL) pyrachlophos;	(CLXXXII) RH-2485;	
(CLI) pyrada-phenthion;	(CLXXXIII) D 2341;	
(CLII) pyresmethrin;	(CLXXXIV) XMC (3,5,-	
(CLIII) pyrethrum;	xylyl methyl-	
(CLIV) RH 5992;	carbamate),	
(CLV) salithion;	(CLXXXV) lufenuron	
(CLVI) sebufos;	(CLXXXVI) fluazuron	
(CLVII) sulfotep;	(CLXXXVII) methoprene	
(CLVIII) sulprofos;	(CLXXXVIII) hydroprene	

Non-limiting examples of suitable anthelmintics are mentioned below, some examples, in addition to having the anthelmintic activity, also having an insecticidal and acaricidal activity and, in some cases, being already contained in the list above:

(A1) praziquantel = 2-cyclohexylcarbonyl-4-oxo-1,2,3,6,7,11b-hexahydro-4H-pyrazino[2,1- α]isoquinoline

(A2) closantel = 3,5-diiodo-N-[5-chloro-2-methyl-4-(α -cyano-4-chlorobenzyl)phenyl]salicyl-amide

(A3) triclabendazole = 5-chloro-6-(2,3-dichlorophenoxy)-2-methylthio-1H-benzimidazole

(A4) levamisole = L-(-)-2,3,5,6-tetrahydro-6-phenylimidazo[2,1b]thiazole

(A5) mebendazole = (5-benzoyl-1H-benzimidazol-2-yl)carbamic acid methyl ester

(A6) omphalotin = a macrocyclic fermentation product of the fungus *Omphalotus olearius* described in WO 97/20857

(A7) abamectin = avermectin B1

(A8) ivermectin = 22,23-dihydroavermectin B1

(A9) moxidectin = 5-O-demethyl-28-deoxy-25-(1,3-dimethyl-1-butenyl)-6,28-epoxy-23-(methoxyimino)-milbemycin B

(A10) doramectin = 25-cyclohexyl-5-O-demethyl-25-de(1-methylpropyl)-avermectin A1a

(A11) milbemectin = mixture of milbemycin A3 and milbemycin A4

(A12) milbemycinoxim = 5-oxime of milbemectin

Non-limiting examples of suitable repelling substances (repellents or detachers) are:

(R1) DEET (N,N-diethyl-m-toluamide)

(R2) KBR 3023 N-butyl-2-oxycarbonyl-(2-hydroxy)-piperidine

(R3) cymiazole = N,-2,3-dihydro-3-methyl-1,3-thiazol-2-ylidene-2,4-xylidene

The said mixing partners are well known to experts in the field. Most of them are described in the various editions of the Pesticide Manual, The British Crop Protection Council, London, and others are described in the various editions of The Merck Index, Merck & Co., Inc., Rahway, New Jersey, USA or in the patent literature. The following list therefore confines itself to giving some examples of sources.

(I) 2-methyl-2-(methylthio)propionaldehyde-O-methylcarbamoyloxime (aldicarb), from The Pesticide Manual, 11th ed. (1997), The British Crop Protection Council, London, page 26;

(II) S-(3,4-dihydro-4-oxobenzo[d]-[1,2,3]-triazin-3-ylmethyl)O,O-dimethyl-phosphorodithioate (azinphos-methyl), from The Pesticide Manual, 11th ed. (1997), The British Crop Protection Council, London, page 67;

- (III) ethyl-N-[2,3-dihydro-2,2-dimethylbenzofuran-7-yloxycarbonyl-(methyl)aminothio]-N-isopropyl- β -alaninate (benfuracarb), from The Pesticide Manual, 11th ed. (1997), The British Crop Protection Council, London, page 96;
- (IV) 2-methylbiphenyl-3-ylmethyl-(Z)-(1*RS*)-*cis*-3-(2-chloro-3,3,3-trifluoroprop-1-enyl)-2,2-dimethylcyclopropanecarboxylate (bifenthrin), from The Pesticide Manual, 11th ed. (1997), The British Crop Protection Council, London, page 118;
- (V) 2-tert-butylimino-3-isopropyl-5-phenyl-1,3,5-thiadiazian-4-one (buprofezin), from The Pesticide Manual, 11th ed. (1997), The British Crop Protection Council, London, page 157;
- (VI) 2,3-dihydro-2,2-dimethylbenzofuran-7-yl-methylcarbamate (carbofuran), from The Pesticide Manual, 11th ed. (1997), The British Crop Protection Council, London, page 186;
- (VII) 2,3-dihydro-2,2-dimethylbenzofuran-7-yl-(dibutylaminothio)methylcarbamate (carbosulfan), from The Pesticide Manual, 11th ed. (1997), The British Crop Protection Council, London, page 188;
- (VIII) *S,S'*-(2-dimethylaminotrimethylene)-bis(thiocarbamate) (cartap), from The Pesticide Manual, 11th ed. (1997), The British Crop Protection Council, London, page 193;
- (IX) 1-[3,5-dichloro-4-(3-chloro-5-trifluoromethyl-2-pyridyloxy)phenyl]-3-(2,6-difluorobenzoyl)-urea (chlorfluazuron), from The Pesticide Manual, 11th ed. (1997), The British Crop Protection Council, London, page 213;
- (X) *O,O*-diethyl-*O*-3,5,6-trichloro-2-pyridyl-phosphorothioate (chlorpyrifos), from The Pesticide Manual, 11th ed. (1997), The British Crop Protection Council, London, page 235;
- (XI) (*RS*)- α -cyano-4-fluoro-3-phenoxybenzyl-(1*RS*,3*RS*;1*RS*,3*RS*)-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate (cyfluthrin), from The Pesticide Manual, 11th ed. (1997), The British Crop Protection Council, London, page 293;
- (XII) mixture of (*S*)- α -cyano-3-phenoxybenzyl-(Z)-(1*R*,3*R*)-3-(2-chloro-3,3,3-trifluoropropenyl)-2,2-dimethylcyclopropanecarboxylate and (*R*)- α -cyano-3-phenoxybenzyl-(Z)-(1*R*,3*R*)-3-(2-chloro-3,3,3-trifluoropropenyl)-2,2-dimethylcyclopropanecarboxylate (lambda-cyhalothrin), from The Pesticide Manual, 11th ed. (1997), The British Crop Protection Council, London, page 300;
- (XIII) racemate consisting of (*S*)- α -cyano-3-phenoxybenzyl-(1*R*,3*R*)-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate and (*R*)- α -cyano-3-phenoxybenzyl-(1*S*,3*S*)-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate (alpha-cypermethrin), from The Pesticide Manual, 11th ed. (1997), The British Crop Protection Council, London, page 308;

- (XIV) a mixture of the stereoisomers of (*S*)- α -cyano-3-phenoxybenzyl (1*RS*,3*RS*,1*RS*,3*RS*)-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate (zeta-cypermethrin), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 314;
- (XV) (*S*)- α -cyano-3-phenoxybenzyl-(1*R*,3*R*)-3-(2,2-dibromovinyl)-2,2-dimethylcyclopropanecarboxylate (deltamethrin), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 344;
- (XVI) (4-chlorophenyl)-3-(2,6-difluorobenzoyl)urea (diflubenzuron), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 395;
- (XVII) (1,4,5,6,7,7-hexachloro-8,9,10-trinorborn-5-en-2,3-ylenebismethylene)-sulfite (endosulfan), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 459;
- (XVIII) α -ethylthio-o-tolyl-methylcarbamate (ethiofencarb), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 479;
- (XIX) *O,O*-dimethyl-*O*-4-nitro-*m*-tolyl-phosphorothioate (fenitrothion), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 514;
- (XX) 2-sec-butylphenyl-methylcarbamate (fenobucarb), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 516;
- (XXI) (*RS*)- α -cyano-3-phenoxybenzyl-(*RS*)-2-(4-chlorophenyl)-3-methylbutyrate (fenvalerate), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 539;
- (XXII) *S*-[formyl(methyl)carbamoylmethyl]-*O,O*-dimethyl-phosphorodithioate (formothion), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 625;
- (XXIII) 4-methylthio-3,5-xylyl-methylcarbamate (methiocarb), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 813;
- (XXIV) 7-chlorobicyclo[3.2.0]hepta-2,6-dien-6-yl-dimethylphosphate (heptenophos), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 670;
- (XXV) 1-(6-chloro-3-pyridylmethyl)-*N*-nitroimidazolidin-2-ylideneamine (imidacloprid), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 706;
- (XXVI) 2-isopropylphenyl-methylcarbamate (isoprocab), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 729;

- (XXVII) *O,S*-dimethyl-phosphoramidothioate (methamidophos), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 808;
- (XXVIII) *S*-methyl-*N*-(methylcarbamoyloxy)thioacetimidate (methomyl), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 815;
- (XXIX) methyl-3-(dimethoxyphosphinoyloxy)but-2-enoate (mevinphos), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 844;
- (XXX) *O,O*-diethyl-*O*-4-nitrophenyl-phosphorothioate (parathion), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 926;
- (XXXI) *O,O*-dimethyl-*O*-4-nitrophenyl-phosphorothioate (parathion-methyl), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 928;
- (XXXII) *S*-6-chloro-2,3-dihydro-2-oxo-1,3-benzoxazol-3-ylmethyl-*O,O*-diethyl-phosphorodithioate (phosalone), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 963;
- (XXXIII) 2-dimethylamino-5,6-dimethylpyrimidin-4-yl-dimethylcarbamate (pirimicarb), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 985;
- (XXXIV) 2-isopropoxyphenyl-methylcarbamate (propoxur), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 1036;
- (XXXV) 1-(3,5-dichloro-2,4-difluorophenyl)-3-(2,6-difluorobenzoyl)urea (teflubenzuron), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 1158;
- (XXXVI) *S*-tert-butylthiomethyl-*O,O*-dimethyl-phosphorodithioate (terbufos), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 1165;
- (XXXVII) ethyl-(3-tert-butyl-1-dimethylcarbamoyl-1*H*-1,2,4-triazol-5-yl-thio)-acetate, (triazamate), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 1224;
- (XXXVIII) abamectin, from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 3;
- (XXXIX) 2-sec-butylphenyl-methylcarbamate (fenobucarb), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 516;
- (XL) *N*-tert-butyl-*N*-(4-ethylbenzoyl)-3,5-dimethylbenzohydrazide (tebufenozide), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 1147;

- (XLI) (±)-5-amino-1-(2,6-dichloro- α,α,α -trifluoro-p-tolyl)-4-trifluoromethyl-sulfinylpyrazole-3-carbonitrile (fipronil), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 545;
- (XLII) (*RS*)- α -cyano-4-fluoro-3-phenoxybenzyl(1*RS*,3*RS*;1*RS*,3*RS*)-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate (beta-cyfluthrin), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 295;
- (XLIII) (4-ethoxyphenyl)-[3-(4-fluoro-3-phenoxyphenyl)propyl](dimethyl)silane (silafluofen), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 1105;
- (XLIV) *tert*-butyl (E)- α -(1,3-dimethyl-5-phenoxy-pyrazol-4-yl-methyleneamino-oxy)-p-toluate (fenpyroximate), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 530;
- (XLV) 2-*tert*-butyl-5-(4-*tert*-butylbenzylthio)-4-chloropyridazin-3(2*H*)-one (pyridaben), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 1161;
- (XLVI) 4-[[4-(1,1-dimethylphenyl)phenyl]ethoxy]-quinazoline (fenazaquin), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 507;
- (XLVII) 4-phenoxyphenyl-(*RS*)-2-(pyridyloxy)propyl ether (pyriproxyfen), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 1073;
- (XLVIII) 5-chloro-*N*-(2-[4-(2-ethoxyethyl)-2,3-dimethylphenoxy]ethyl)-6-ethylpyrimidine-4-amine (pyrimidifen), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 1070;
- (XLIX) (E)-*N*-(6-chloro-3-pyridylmethyl)-*N*-ethyl-*N*-methyl-2-nitrovinylidenediamine (nitenpyram), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 880;
- (L) (E)-*N*¹-[(6-chloro-3-pyridyl)methyl]-*N*²-cyano-*N*¹-methylacetamidine (NI-25, acetamiprid), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 9;
- (LI) avermectin B₁, from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 3;
- (LII) an insect-active extract from a plant, especially (2*R*,6*aS*,12*aS*)-1,2,6,6*a*,12,12*a*-hexahydro-2-isopropenyl-8,9-dimethoxy-chromeno[3,4-*b*]furo[2,3-*h*]chromen-6-one (rotenone), from The Pesticide Manual, 11thed. (1997), The British Crop Protection

- Council, London, page 1097; and an extract from *Azadirachta indica*, especially azadirachtin, from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 59; and
- (LIII) a preparation containing insect-active nematodes, preferably *Heterorhabditis bacteriophora* and *Heterorhabditis megidis*, from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 671; *Steinernema feltiae*, from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 1115, and *Steinernema scapterisci*, from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 1116;
- (LIV) a preparation, obtainable from *Bacillus subtilis*, from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 72; or from a *Bacillus thuringiensis* strain with the exception of compounds isolated from GC91 or from NCTC11821; The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 73;
- (LV) a preparation containing insect-active fungi, preferably *Verticillium lecanii*, from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 1266; *Beauveria brogniartii*, from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 85; and *Beauveria bassiana*, from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 83;
- (LVI) a preparation containing insect-active viruses, preferably *Neodiprion Sertifer NPV*, from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 1342; *Mamestra brassicae NPV*, from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 759; and *Cydia pomonella granulosus* virus, from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 291;
- (CLXXXI) 7-chloro-2,3,4a,5-tetrahydro-2-[methoxycarbonyl(4-trifluoromethoxyphenyl)carbamoyl]indol[1,2e]oxazoline-4a-carboxylate (DPX-MP062, indoxycarb), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 453;
- (CLXXXII) *N'*-tert-butyl-*N'*-(3,5-dimethylbenzoyl)-3-methoxy-2-methylbenzohydrazide (RH-2485, methoxyfenozide), from The Pesticide Manual, 11thed. (1997), The British Crop Protection Council, London, page 1094; and
- (CLXXXIII) (*N'*-[4-methoxy-biphenyl-3-yl]-hydrazinecarboxylic acid isopropyl ester (D 2341), from Brighton Crop Protection Conference, 1996, 487- 493;

(R2) Book of Abstracts, 212th ACS National Meeting Orlando, FL, August 25-29 (1996),

AGRO-020. Publisher: American Chemical Society, Washington, D.C. CONEN: 63BFAF.

In accordance with the above remarks, a further important aspect of the present invention relates to combination preparations for controlling parasites on warm-blooded animals which comprise, in addition to a compound of formula (I), at least one further active ingredient having the same or a different direction of action and at least one physiologically tolerable carrier. The present invention is not restricted to two-component combinations.

Within the scope of the present invention preference is given, for example, to the following two-component combinations, the figure in parenthesis representing one of the combination partners mentioned above and the number following the symbol "&" representing a compound from the Tables which follow:

(IX) & 1.001; (IX) & 1.008; (IX) & 1.011; (IX) & 1.012; (IX) & 1.013; (IX) & 1.014; (IX) & 1.015; (IX) & 1.018; (IX) & 1.019; (IX) & 1.020; (IX) & 1.021; (IX) & 1.022; (IX) & 1.054; (IX) & 1.055; (IX) & 1.056; (IX) & 1.057; (IX) & 1.058; (XIII) & 1.001; (XIII) & 1.008; (XIII) & 1.011; (XIII) & 1.012; (XIII) & 1.013; (XIII) & 1.014; (XIII) & 1.015; (XIII) & 1.018; (XIII) & 1.019; (XIII) & 1.020; (XIII) & 1.021; (XIII) & 1.022; (XIII) & 1.054; (XIII) & 1.055; (XIII) & 1.056; (XIII) & 1.057; (XIII) & 1.058; (XIV) & 1.001; (XIV) & 1.008; (XIV) & 1.011; (XIV) & 1.012; (XIV) & 1.013; (XIV) & 1.014; (XIV) & 1.015; (XIV) & 1.018; (XIV) & 1.019; (XIV) & 1.020; (XIV) & 1.021; (XIV) & 1.022; (XIV) & 1.054; (XIV) & 1.055; (XIV) & 1.056; (XIV) & 1.057; (XIV) & 1.058; (LI) & 1.001; (LI) & 1.008; (LI) & 1.011; (LI) & 1.012; (LI) & 1.013; (LI) & 1.014; (LI) & 1.015; (LI) & 1.018; (LI) & 1.019; (LI) & 1.020; (LI) & 1.021; (LI) & 1.022; (LI) & 1.054; (LI) & 1.055; (LI) & 1.056; (LI) & 1.057; (LI) & 1.058; (XXV) & 1.001; (XXV) & 1.008; (XXV) & 1.011; (XXV) & 1.012; (XXV) & 1.013; (XXV) & 1.014; (XXV) & 1.015; (XXV) & 1.018; (XXV) & 1.019; (XXV) & 1.020; (XXV) & 1.021; (XXV) & 1.022; (XXV) & 1.054; (XXV) & 1.055; (XXV) & 1.056; (XXV) & 1.057; (XXV) & 1.058; (XXXV) & 1.001; (XXXV) & 1.008; (XXXV) & 1.011; (XXXV) & 1.012; (XXXV) & 1.013; (XXXV) & 1.014; (XXXV) & 1.015; (XXXV) & 1.018; (XXXV) & 1.019; (XXXV) & 1.020; (XXXV) & 1.021; (XXXV) & 1.022; (XXXV) & 1.054; (XXXV) & 1.055; (XXXV) & 1.056; (XXXV) & 1.057; (XXXV) & 1.058; (XXXVIII) & 1.001; (XXXVIII) & 1.008; (XXXVIII) & 1.011; (XXXVIII) & 1.012; (XXXVIII) & 1.013; (XXXVIII) & 1.014; (XXXVIII) & 1.015; (XXXVIII) & 1.018; (XXXVIII) & 1.019; (XXXVIII) & 1.020; (XXXVIII) & 1.021; (XXXVIII) & 1.022; (XXXVIII) & 1.054; (XXXVIII) & 1.055;

(XXXVIII) & 1.056; (XXXVIII) & 1.057; (XXXVIII) & 1.058; (XLI) & 1.001; (XLI) & 1.008; (XLI) & 1.011; (XLI) & 1.012; (XLI) & 1.013; (XLI) & 1.014; (XLI) & 1.015; (XLI) & 1.018; (XLI) & 1.019; (XLI) & 1.020; (XLI) & 1.021; (XLI) & 1.022; (XLI) & 1.054; (XLI) & 1.055; (XLI) & 1.056; (XLI) & 1.057; (XLI) & 1.058; (XLVII) & 1.001; (XLVII) & 1.008; (XLVII) & 1.011; (XLVII) & 1.012; (XLVII) & 1.013; (XLVII) & 1.014; (XLVII) & 1.015; (XLVII) & 1.018; (XLVII) & 1.019; (XLVII) & 1.020; (XLVII) & 1.021; (XLVII) & 1.022; (XLVII) & 1.054; (XLVII) & 1.055; (XLVII) & 1.056; (XLVII) & 1.057; (XLVII) & 1.058; (XLIX) & 1.001; (XLIX) & 1.008; (XLIX) & 1.011; (XLIX) & 1.012; (XLIX) & 1.013; (XLIX) & 1.014; (XLIX) & 1.015; (XLIX) & 1.018; (XLIX) & 1.019; (XLIX) & 1.020; (XLIX) & 1.021; (XLIX) & 1.022; (XLIX) & 1.054; (XLIX) & 1.055; (XLIX) & 1.056; (XLIX) & 1.057; (XLIX) & 1.058; (LXI) & 1.001; (LXI) & 1.008; (LXI) & 1.011; (LXI) & 1.012; (LXI) & 1.013; (LXI) & 1.014; (LXI) & 1.015; (LXI) & 1.018; (LXI) & 1.019; (LXI) & 1.020; (LXI) & 1.021; (LXI) & 1.022; (LXI) & 1.054; (LXI) & 1.055; (LXI) & 1.056; (LXI) & 1.057; (LXI) & 1.058; (LXII) & 1.001; (LXII) & 1.008; (LXII) & 1.011; (LXII) & 1.012; (LXII) & 1.013; (LXII) & 1.014; (LXII) & 1.015; (LXII) & 1.018; (LXII) & 1.019; (LXII) & 1.020; (LXII) & 1.021; (LXII) & 1.022; (LXII) & 1.054; (LXII) & 1.055; (LXII) & 1.056; (LXII) & 1.057; (LXII) & 1.058; (CIX) & 1.001; (CIX) & 1.008; (CIX) & 1.011; (CIX) & 1.012; (CIX) & 1.013; (CIX) & 1.014; (CIX) & 1.015; (CIX) & 1.018; (CIX) & 1.019; (CIX) & 1.020; (CIX) & 1.021; (CIX) & 1.022; (CIX) & 1.054; (CIX) & 1.055; (CIX) & 1.056; (CIX) & 1.057; (CIX) & 1.058; (CXIII) & 1.001; (CXIII) & 1.008; (CXIII) & 1.011; (CXIII) & 1.012; (CXIII) & 1.013; (CXIII) & 1.014; (CXIII) & 1.015; (CXIII) & 1.018; (CXIII) & 1.019; (CXIII) & 1.020; (CXIII) & 1.021; (CXIII) & 1.022; (CXIII) & 1.054; (CXIII) & 1.055; (CXIII) & 1.056; (CXIII) & 1.057; (CXIII) & 1.058; (CXIX) & 1.001; (CXIX) & 1.008; (CXIX) & 1.011; (CXIX) & 1.012; (CXIX) & 1.013; (CXIX) & 1.014; (CXIX) & 1.015; (CXIX) & 1.018; (CXIX) & 1.019; (CXIX) & 1.020; (CXIX) & 1.021; (CXIX) & 1.022; (CXIX) & 1.054; (CXIX) & 1.055; (CXIX) & 1.056; (CXIX) & 1.057; (CXIX) & 1.058; (CXXIV) & 1.001; (CXXIV) & 1.008; (CXXIV) & 1.011; (CXXIV) & 1.012; (CXXIV) & 1.013; (CXXIV) & 1.014; (CXXIV) & 1.015; (CXXIV) & 1.018; (CXXIV) & 1.019; (CXXIV) & 1.020; (CXXIV) & 1.021; (CXXIV) & 1.022; (CXXIV) & 1.054; (CXXIV) & 1.055; (CXXIV) & 1.056; (CXXIV) & 1.057; (CXXIV) & 1.058; (CXXXI) & 1.001; (CXXXI) & 1.008; (CXXXI) & 1.011; (CXXXI) & 1.012; (CXXXI) & 1.013; (CXXXI) & 1.014; (CXXXI) & 1.015; (CXXXI) & 1.018; (CXXXI) & 1.019; (CXXXI) & 1.020; (CXXXI) & 1.021; (CXXXI) & 1.022; (CXXXI) & 1.054; (CXXXI) & 1.055; (CXXXI) & 1.056; (CXXXI) & 1.057; (CXXXI) & 1.058; (CXXXII) & 1.001; (CXXXII) & 1.008; (CXXXII) & 1.011; (CXXXII) & 1.012; (CXXXII) & 1.013; (CXXXII) & 1.014; (CXXXII) & 1.015; (CXXXII) & 1.018; (CXXXII) & 1.019; (CXXXII) & 1.020; (CXXXII) & 1.021; (CXXXII) & 1.022; (CXXXII) & 1.054; (CXXXII) & 1.055; (CXXXII) &

1.056; (CXXXII) & 1.057; (CXXXII) & 1.058; (CXXXIX) & 1.001; (CXXXIX) & 1.008; (CXXXIX) & 1.011; (CXXXIX) & 1.012; (CXXXIX) & 1.013; (CXXXIX) & 1.014; (CXXXIX) & 1.015; (CXXXIX) & 1.018; (CXXXIX) & 1.019; (CXXXIX) & 1.020; (CXXXIX) & 1.021; (CXXXIX) & 1.022; (CXXXIX) & 1.054; (CXXXIX) & 1.055; (CXXXIX) & 1.056; (CXXXIX) & 1.057; (CXXXIX) & 1.058; (CLII) & 1.001; (CLII) & 1.008; (CLII) & 1.011; (CLII) & 1.012; (CLII) & 1.013; (CLII) & 1.014; (CLII) & 1.015; (CLII) & 1.018; (CLII) & 1.019; (CLII) & 1.020; (CLII) & 1.021; (CLII) & 1.022; (CLII) & 1.054; (CLII) & 1.055; (CLII) & 1.056; (CLII) & 1.057; (CLII) & 1.058; (CLIII) & 1.001; (CLIII) & 1.008; (CLIII) & 1.011; (CLIII) & 1.012; (CLIII) & 1.013; (CLIII) & 1.014; (CLIII) & 1.015; (CLIII) & 1.018; (CLIII) & 1.019; (CLIII) & 1.020; (CLIII) & 1.021; (CLIII) & 1.022; (CLIII) & 1.054; (CLIII) & 1.055; (CLIII) & 1.056; (CLIII) & 1.057; (CLIII) & 1.058; (CLIX) & 1.001; (CLIX) & 1.008; (CLIX) & 1.011; (CLIX) & 1.012; (CLIX) & 1.013; (CLIX) & 1.014; (CLIX) & 1.015; (CLIX) & 1.018; (CLIX) & 1.019; (CLIX) & 1.020; (CLIX) & 1.021; (CLIX) & 1.022; (CLIX) & 1.054; (CLIX) & 1.055; (CLIX) & 1.056; (CLIX) & 1.057; (CLIX) & 1.058; (CLXXIII) & 1.001; (CLXXIII) & 1.008; (CLXXIII) & 1.011; (CLXXIII) & 1.012; (CLXXIII) & 1.013; (CLXXIII) & 1.014; (CLXXIII) & 1.015; (CLXXIII) & 1.018; (CLXXIII) & 1.019; (CLXXIII) & 1.020; (CLXXIII) & 1.021; (CLXXIII) & 1.022; (CLXXIII) & 1.054; (CLXXIII) & 1.055; (CLXXIII) & 1.056; (CLXXIII) & 1.057; (CLXXIII) & 1.058; (CLXXV) & 1.001; (CLXXV) & 1.008; (CLXXV) & 1.011; (CLXXV) & 1.012; (CLXXV) & 1.013; (CLXXV) & 1.014; (CLXXV) & 1.015; (CLXXV) & 1.018; (CLXXV) & 1.019; (CLXXV) & 1.020; (CLXXV) & 1.021; (CLXXV) & 1.022; (CLXXV) & 1.054; (CLXXV) & 1.055; (CLXXV) & 1.056; (CLXXV) & 1.057; (CLXXV) & 1.058; (CLXXXI) & 1.001; (CLXXXI) & 1.008; (CLXXXI) & 1.011; (CLXXXI) & 1.012; (CLXXXI) & 1.013; (CLXXXI) & 1.014; (CLXXXI) & 1.015; (CLXXXI) & 1.018; (CLXXXI) & 1.019; (CLXXXI) & 1.020; (CLXXXI) & 1.021; (CLXXXI) & 1.022; (CLXXXI) & 1.054; (CLXXXI) & 1.055; (CLXXXI) & 1.056; (CLXXXI) & 1.057; (CLXXXI) & 1.058; (CLXXXV) & 1.001; (CLXXXV) & 1.008; (CLXXXV) & 1.011; (CLXXXV) & 1.012; (CLXXXV) & 1.013; (CLXXXV) & 1.014; (CLXXXV) & 1.015; (CLXXXV) & 1.018; (CLXXXV) & 1.019; (CLXXXV) & 1.020; (CLXXXV) & 1.021; (CLXXXV) & 1.022; (CLXXXV) & 1.054; (CLXXXV) & 1.055; (CLXXXV) & 1.056; (CLXXXV) & 1.057; (CLXXXV) & 1.058; (CLXXXVI) & 1.001; (CLXXXVI) & 1.008; (CLXXXVI) & 1.011; (CLXXXVI) & 1.012; (CLXXXVI) & 1.013; (CLXXXVI) & 1.014; (CLXXXVI) & 1.015; (CLXXXVI) & 1.018; (CLXXXVI) & 1.019; (CLXXXVI) & 1.020; (CLXXXVI) & 1.021; (CLXXXVI) & 1.022; (CLXXXVI) & 1.054; (CLXXXVI) & 1.055; (CLXXXVI) & 1.056; (CLXXXVI) & 1.057; (CLXXXVI) & 1.058; (CLXXXVII) & 1.001; (CLXXXVII) & 1.008; (CLXXXVII) & 1.011; (CLXXXVII) & 1.012; (CLXXXVII) & 1.013; (CLXXXVII) & 1.014;

(CLXXXVII) & 1.015; (CLXXXVII) & 1.018; (CLXXXVII) & 1.019; (CLXXXVII) & 1.020;
(CLXXXVII) & 1.021; (CLXXXVII) & 1.022; (CLXXXVII) & 1.054; (CLXXXVII) & 1.055;
(CLXXXVII) & 1.056; (CLXXXVII) & 1.057; (CLXXXVII) & 1.058; (CLXXXVIII) & 1.001;
(CLXXXVIII) & 1.008; (CLXXXVIII) & 1.011; (CLXXXVIII) & 1.012; (CLXXXVIII) & 1.013;
(CLXXXVIII) & 1.014; (CLXXXVIII) & 1.015; (CLXXXVIII) & 1.018; (CLXXXVIII) & 1.019;
(CLXXXVIII) & 1.020; (CLXXXVIII) & 1.021; (CLXXXVIII) & 1.022; (CLXXXVIII) & 1.054;
(CLXXXVIII) & 1.055; (CLXXXVIII) & 1.056; (CLXXXVIII) & 1.057; (CLXXXVIII) & 1.058;
(CLXXXIX) & 1.001; (CLXXXIX) & 1.008; (CLXXXIX) & 1.011; (CLXXXIX) & 1.012;
(CLXXXIX) & 1.013; (CLXXXIX) & 1.014; (CLXXXIX) & 1.015; (CLXXXIX) & 1.018;
(CLXXXIX) & 1.019; (CLXXXIX) & 1.020; (CLXXXIX) & 1.021; (CLXXXIX) & 1.022;
(CLXXXIX) & 1.054; (CLXXXIX) & 1.055; (CLXXXIX) & 1.056; (CLXXXIX) & 1.057;
(CLXXXIX) & 1.058; (CXC) & 1.001; (CXC) & 1.008; (CXC) & 1.011; (CXC) & 1.012; (CXC)
& 1.013; (CXC) & 1.014; (CXC) & 1.015; (CXC) & 1.018; (CXC) & 1.019; (CXC) & 1.020;
(CXC) & 1.021; (CXC) & 1.022; (CXC) & 1.054; (CXC) & 1.055; (CXC) & 1.056; (CXC) &
1.057; (CXC) & 1.058; (CXCL) & 1.001; (CXCL) & 1.008; (CXCL) & 1.011; (CXCL) & 1.012;
(CXCL) & 1.013; (CXCL) & 1.014; (CXCL) & 1.015; (CXCL) & 1.018; (CXCL) & 1.019;
(CXCL) & 1.020; (CXCL) & 1.021; (CXCL) & 1.022; (CXCL) & 1.054; (CXCL) & 1.055;
(CXCL) & 1.056; (CXCL) & 1.057; (CXCL) & 1.058; (A1) & 1.001; (A1) & 1.008; (A1) &
1.011; (A1) & 1.012; (A1) & 1.013; (A1) & 1.014; (A1) & 1.015; (A1) & 1.018; (A1) & 1.019;
(A1) & 1.020; (A1) & 1.021; (A1) & 1.022; (A1) & 1.054; (A1) & 1.055; (A1) & 1.056; (A1) &
1.057; (A1) & 1.058; (A2) & 1.001; (A2) & 1.008; (A2) & 1.011; (A2) & 1.012; (A2) & 1.013;
(A2) & 1.014; (A2) & 1.015; (A2) & 1.018; (A2) & 1.019; (A2) & 1.020; (A2) & 1.021; (A2) &
1.022; (A2) & 1.054; (A2) & 1.055; (A2) & 1.056; (A2) & 1.057; (A2) & 1.058; (A3) & 1.001;
(A3) & 1.008; (A3) & 1.011; (A3) & 1.012; (A3) & 1.013; (A3) & 1.014; (A3) & 1.015; (A3) &
1.018; (A3) & 1.019; (A3) & 1.020; (A3) & 1.021; (A3) & 1.022; (A3) & 1.054; (A3) & 1.055;
(A3) & 1.056; (A3) & 1.057; (A3) & 1.058; (A4) & 1.001; (A4) & 1.008; (A4) & 1.011; (A4) &
1.012; (A4) & 1.013; (A4) & 1.014; (A4) & 1.015; (A4) & 1.018; (A4) & 1.019; (A4) & 1.020;
(A4) & 1.021; (A4) & 1.022; (A4) & 1.054; (A4) & 1.055; (A4) & 1.056; (A4) & 1.057; (A4) &
1.058; (A5) & 1.001; (A5) & 1.008; (A5) & 1.011; (A5) & 1.012; (A5) & 1.013; (A5) & 1.014;
(A5) & 1.015; (A5) & 1.018; (A5) & 1.019; (A5) & 1.020; (A5) & 1.021; (A5) & 1.022; (A5) &
1.054; (A5) & 1.055; (A5) & 1.056; (A5) & 1.057; (A5) & 1.058; (A6) & 1.001; (A6) & 1.008;
(A6) & 1.011; (A6) & 1.012; (A6) & 1.013; (A6) & 1.014; (A6) & 1.015; (A6) & 1.018; (A6) &
1.019; (A6) & 1.020; (A6) & 1.021; (A6) & 1.022; (A6) & 1.054; (A6) & 1.055; (A6) & 1.056;
(A6) & 1.057; (A6) & 1.058; (A10) & 1.001; (A10) & 1.008; (A10) & 1.011; (A10) & 1.012;


(A10) & 1.013; (A10) & 1.014; (A10) & 1.015; (A10) & 1.018; (A10) & 1.019; (A10) & 1.020;
(A10) & 1.021; (A10) & 1.022; (A10) & 1.054; (A10) & 1.055; (A10) & 1.056; (A10) & 1.057;
(A10) & 1.058; (A11) & 1.001; (A11) & 1.008; (A11) & 1.011; (A11) & 1.012; (A11) & 1.013;
(A11) & 1.014; (A11) & 1.015; (A11) & 1.018; (A11) & 1.019; (A11) & 1.020; (A11) & 1.021;
(A11) & 1.022; (A11) & 1.054; (A11) & 1.055; (A11) & 1.056; (A11) & 1.057; (A11) & 1.058;
(R1) & 1.001; (R1) & 1.008; (R1) & 1.011; (R1) & 1.012; (R1) & 1.013; (R1) & 1.014; (R1) &
1.015; (R1) & 1.018; (R1) & 1.019; (R1) & 1.020; (R1) & 1.021; (R1) & 1.022; (R1) & 1.054;
(R1) & 1.055; (R1) & 1.056; (R1) & 1.057; (R1) & 1.058; (R2) & 1.001; (R2) & 1.008; (R2) &
1.011; (R2) & 1.012; (R2) & 1.013; (R2) & 1.014; (R2) & 1.015; (R2) & 1.018; (R2) & 1.019;
(R2) & 1.020; (R2) & 1.021; (R2) & 1.022; (R2) & 1.054; (R2) & 1.055; (R2) & 1.056; (R2) &
1.057; (R2) & 1.058; (R3) & 1.001; (R3) & 1.008; (R3) & 1.011; (R3) & 1.012; (R3) & 1.013;
(R3) & 1.014; (R3) & 1.015; (R3) & 1.018; (R3) & 1.019; (R3) & 1.020; (R3) & 1.021; (R3) &
1.022; (R3) & 1.054; (R3) & 1.055; (R3) & 1.056; (R3) & 1.057; (R3) & 1.058; (IX) & 1.059;
(IX) & 1.060; (IX) & 1.061; (IX) & 1.062; (IX) & 1.063; (IX) & 1.064; (IX) & 1.065; (IX) &
1.066; (IX) & 1.067; (IX) & 1.068; (IX) & 1.069; (IX) & 1.070; (IX) & 1.071; (IX) & 1.072; (IX)
& 1.073; (IX) & 1.074; (IX) & 1.075; (IX) & 1.076; (IX) & 1.077; (IX) & 1.078; (IX) & 1.079;
(IX) & 1.080; (IX) & 1.081; (IX) & 1.082; (IX) & 1.083; (IX) & 1.084; (IX) & 1.085; (IX) &
1.086; (IX) & 1.087; (XIII) & 1.059; (XIII) & 1.060; (XIII) & 1.061; (XIII) & 1.062; (XIII) & 1.063;
(XIII) & 1.064; (XIII) & 1.065; (XIII) & 1.066; (XIII) & 1.067; (XIII) & 1.068; (XIII) & 1.069; (XIII)
& 1.070; (XIII) & 1.071; (XIII) & 1.072; (XIII) & 1.073; (XIII) & 1.074; (XIII) & 1.075; (XIII) &
1.076; (XIII) & 1.077; (XIII) & 1.078; (XIII) & 1.079; (XIII) & 1.080; (XIII) & 1.081; (XIII) &
1.082; (XIII) & 1.083; (XIII) & 1.084; (XIII) & 1.085; (XIII) & 1.086; (XIII) & 1.087; (XIV) &
1.059; (XIV) & 1.060; (XIV) & 1.061; (XIV) & 1.062; (XIV) & 1.063; (XIV) & 1.064; (XIV) &
1.065; (XIV) & 1.066; (XIV) & 1.067; (XIV) & 1.068; (XIV) & 1.069; (XIV) & 1.070; (XIV) &
1.071; (XIV) & 1.072; (XIV) & 1.073; (XIV) & 1.074; (XIV) & 1.075; (XIV) & 1.076; (XIV) &
1.077; (XIV) & 1.078; (XIV) & 1.079; (XIV) & 1.080; (XIV) & 1.081; (XIV) & 1.082; (XIV) &
1.083; (XIV) & 1.084; (XIV) & 1.085; (XIV) & 1.086; (XIV) & 1.087; (LI) & 1.059; (LI) & 1.060;
(LI) & 1.061; (LI) & 1.062; (LI) & 1.063; (LI) & 1.064; (LI) & 1.065; (LI) & 1.066; (LI) & 1.067;
(LI) & 1.068; (LI) & 1.069; (LI) & 1.070; (LI) & 1.071; (LI) & 1.072; (LI) & 1.073; (LI) & 1.074;
(LI) & 1.075; (LI) & 1.076; (LI) & 1.077; (LI) & 1.078; (LI) & 1.079; (LI) & 1.080; (LI) & 1.081;
(LI) & 1.082; (LI) & 1.083; (LI) & 1.084; (LI) & 1.085; (LI) & 1.086; (LI) & 1.087; (XXV) &
1.059; (XXV) & 1.060; (XXV) & 1.061; (XXV) & 1.062; (XXV) & 1.063; (XXV) & 1.064; (XXV)
& 1.065; (XXV) & 1.066; (XXV) & 1.067; (XXV) & 1.068; (XXV) & 1.069; (XXV) & 1.070;
(XXV) & 1.071; (XXV) & 1.072; (XXV) & 1.073; (XXV) & 1.074; (XXV) & 1.075; (XXV) &

1.076; (XXV) & 1.077; (XXV) & 1.078; (XXV) & 1.079; (XXV) & 1.080; (XXV) & 1.081; (XXV) & 1.082; (XXV) & 1.083; (XXV) & 1.084; (XXV) & 1.085; (XXV) & 1.086; (XXV) & 1.087; (XXXV) & 1.059; (XXXV) & 1.060; (XXXV) & 1.061; (XXXV) & 1.062; (XXXV) & 1.063; (XXXV) & 1.064; (XXXV) & 1.065; (XXXV) & 1.066; (XXXV) & 1.067; (XXXV) & 1.068; (XXXV) & 1.069; (XXXV) & 1.070; (XXXV) & 1.071; (XXXV) & 1.072; (XXXV) & 1.073; (XXXV) & 1.074; (XXXV) & 1.075; (XXXV) & 1.076; (XXXV) & 1.077; (XXXV) & 1.078; (XXXV) & 1.079; (XXXV) & 1.080; (XXXV) & 1.081; (XXXV) & 1.082; (XXXV) & 1.083; (XXXV) & 1.084; (XXXV) & 1.085; (XXXV) & 1.086; (XXXV) & 1.087; (XXXVIII) & 1.059; (XXXVIII) & 1.060; (XXXVIII) & 1.061; (XXXVIII) & 1.062; (XXXVIII) & 1.063; (XXXVIII) & 1.064; (XXXVIII) & 1.065; (XXXVIII) & 1.066; (XXXVIII) & 1.067; (XXXVIII) & 1.068; (XXXVIII) & 1.069; (XXXVIII) & 1.070; (XXXVIII) & 1.071; (XXXVIII) & 1.072; (XXXVIII) & 1.073; (XXXVIII) & 1.074; (XXXVIII) & 1.075; (XXXVIII) & 1.076; (XXXVIII) & 1.077; (XXXVIII) & 1.078; (XXXVIII) & 1.079; (XXXVIII) & 1.080; (XXXVIII) & 1.081; (XXXVIII) & 1.082; (XXXVIII) & 1.083; (XXXVIII) & 1.084; (XXXVIII) & 1.085; (XXXVIII) & 1.086; (XXXVIII) & 1.087; (XLI) & 1.059; (XLI) & 1.060; (XLI) & 1.061; (XLI) & 1.062; (XLI) & 1.063; (XLI) & 1.064; (XLI) & 1.065; (XLI) & 1.066; (XLI) & 1.067; (XLI) & 1.068; (XLI) & 1.069; (XLI) & 1.070; (XLI) & 1.071; (XLI) & 1.072; (XLI) & 1.073; (XLI) & 1.074; (XLI) & 1.075; (XLI) & 1.076; (XLI) & 1.077; (XLI) & 1.078; (XLI) & 1.079; (XLI) & 1.080; (XLI) & 1.081; (XLI) & 1.082; (XLI) & 1.083; (XLI) & 1.084; (XLI) & 1.085; (XLI) & 1.086; (XLI) & 1.087; (XLVII) & 1.059; (XLVII) & 1.060; (XLVII) & 1.061; (XLVII) & 1.062; (XLVII) & 1.063; (CIX) & 1.064; (XLVII) & 1.065; (XLVII) & 1.066; (XLVII) & 1.067; (XLVII) & 1.068; (XLVII) & 1.069; (XLVII) & 1.070; (XLVII) & 1.071; (XLVII) & 1.072; (XLVII) & 1.073; (XLVII) & 1.074; (XLVII) & 1.075; (XLVII) & 1.076; (XLVII) & 1.077; (XLVII) & 1.078; (XLVII) & 1.079; (XLVII) & 1.080; (XLVII) & 1.081; (XLVII) & 1.082; (XLVII) & 1.083; (XLVII) & 1.084; (XLVII) & 1.085; (XLVII) & 1.086; (XLVII) & 1.087; (XLIX) & 1.059; (XLIX) & 1.060; (XLIX) & 1.061; (XLIX) & 1.062; (XLIX) & 1.063; (CIX) & 1.064; (XLIX) & 1.065; (XLIX) & 1.066; (XLIX) & 1.067; (XLIX) & 1.068; (XLIX) & 1.069; (XLIX) & 1.070; (XLIX) & 1.071; (XLIX) & 1.072; (XLIX) & 1.073; (XLIX) & 1.074; (XLIX) & 1.075; (XLIX) & 1.076; (XLIX) & 1.077; (XLIX) & 1.078; (XLIX) & 1.079; (XLIX) & 1.080; (XLIX) & 1.081; (XLIX) & 1.082; (XLIX) & 1.083; (XLIX) & 1.084; (XLIX) & 1.085; (XLIX) & 1.086; (XLIX) & 1.087; (LXI) & 1.059; (LXI) & 1.060; (LXI) & 1.061; (LXI) & 1.062; (LXI) & 1.063; (CIX) & 1.064; (LXI) & 1.065; (LXI) & 1.066; (LXI) & 1.067; (LXI) & 1.068; (LXI) & 1.069; (LXI) & 1.070; (LXI) & 1.071; (LXI) & 1.072; (LXI) & 1.073; (LXI) & 1.074; (LXI) & 1.075; (LXI) & 1.076; (LXI) & 1.077; (LXI) & 1.078; (LXI) & 1.079; (LXI) & 1.080; (LXI) & 1.081; (LXI) & 1.082; (LXI) & 1.083; (LXI) & 1.084; (LXI) & 1.085;

(LXI) & 1.086; (LXI) & 1.087; (LXII) & 1.059; (LXII) & 1.060; (LXII) & 1.061; (LXII) & 1.062; (LXII) & 1.063; (CIX) & 1.064; (LXII) & 1.065; (LXII) & 1.066; (LXII) & 1.067; (LXII) & 1.068; (LXII) & 1.069; (LXII) & 1.070; (LXII) & 1.071; (LXII) & 1.072; (LXII) & 1.073; (LXII) & 1.074; (LXII) & 1.075; (LXII) & 1.076; (LXII) & 1.077; (LXII) & 1.078; (LXII) & 1.079; (LXII) & 1.080; (LXII) & 1.081; (LXII) & 1.082; (LXII) & 1.083; (LXII) & 1.084; (LXII) & 1.085; (LXII) & 1.086; (LXII) & 1.087; (CIX) & 1.059; (CIX) & 1.060; (CIX) & 1.061; (CIX) & 1.062; (CIX) & 1.063; (CIX) & 1.064; (CIX) & 1.065; (CIX) & 1.066; (CIX) & 1.067; (CIX) & 1.068; (CIX) & 1.069; (CIX) & 1.070; (CIX) & 1.071; (CIX) & 1.072; (CIX) & 1.073; (CIX) & 1.074; (CIX) & 1.075; (CIX) & 1.076; (CIX) & 1.077; (CIX) & 1.078; (CIX) & 1.079; (CIX) & 1.080; (CIX) & 1.081; (CIX) & 1.082; (CIX) & 1.083; (CIX) & 1.084; (CIX) & 1.085; (CIX) & 1.086; (CIX) & 1.087; (CXIII) & 1.059; (CXIII) & 1.060; (CXIII) & 1.061; (CXIII) & 1.062; (CXIII) & 1.063; (CXIII) & 1.064; (CXIII) & 1.065; (CXIII) & 1.066; (CXIII) & 1.067; (CXIII) & 1.068; (CXIII) & 1.069; (CXIII) & 1.070; (CXIII) & 1.071; (CXIII) & 1.072; (CXIII) & 1.073; (CXIII) & 1.074; (CXIII) & 1.075; (CXIII) & 1.076; (CXIII) & 1.077; (CXIII) & 1.078; (CXIII) & 1.079; (CXIII) & 1.080; (CXIII) & 1.081; (CXIII) & 1.082; (CXIII) & 1.083; (CXIII) & 1.084; (CXIII) & 1.085; (CXIII) & 1.086; (CXIII) & 1.087; (CXIX) & 1.059; (CXIX) & 1.060; (CXIX) & 1.061; (CXIX) & 1.062; (CXIX) & 1.063; (CXIX) & 1.064; (CXIX) & 1.065; (CXIX) & 1.066; (CXIX) & 1.067; (CXIX) & 1.068; (CXIX) & 1.069; (CXIX) & 1.070; (CXIX) & 1.071; (CXIX) & 1.072; (CXIX) & 1.073; (CXIX) & 1.074; (CXIX) & 1.075; (CXIX) & 1.076; (CXIX) & 1.077; (CXIX) & 1.078; (CXIX) & 1.079; (CXIX) & 1.080; (CXIX) & 1.081; (CXIX) & 1.082; (CXIX) & 1.083; (CXIX) & 1.084; (CXIX) & 1.085; (CXIX) & 1.086; (CXIX) & 1.087; (CXXIV) & 1.059; (CXXIV) & 1.060; (CXXIV) & 1.061; (CXXIV) & 1.062; (CXXIV) & 1.063; (CXXIV) & 1.064; (CXXIV) & 1.065; (CXXIV) & 1.066; (CXXIV) & 1.067; (CXXIV) & 1.068; (CXXIV) & 1.069; (CXXIV) & 1.070; (CXXIV) & 1.071; (CXXIV) & 1.072; (CXXIV) & 1.073; (CXXIV) & 1.074; (CXXIV) & 1.075; (CXXIV) & 1.076; (CXXIV) & 1.077; (CXXIV) & 1.078; (CXXIV) & 1.079; (CXXIV) & 1.080; (CXXIV) & 1.081; (CXXIV) & 1.082; (CXXIV) & 1.083; (CXXIV) & 1.084; (CXXIV) & 1.085; (CXXIV) & 1.086; (CXXIV) & 1.087; (CXXXI) & 1.059; (CXXXI) & 1.060; (CXXXI) & 1.061; (CXXXI) & 1.062; (CXXXI) & 1.063; (CXXXI) & 1.064; (CXXXI) & 1.065; (CXXXI) & 1.066; (CXXXI) & 1.067; (CXXXI) & 1.068; (CXXXI) & 1.069; (CXXXI) & 1.070; (CXXXI) & 1.071; (CXXXI) & 1.072; (CXXXI) & 1.073; (CXXXI) & 1.074; (CXXXI) & 1.075; (CXXXI) & 1.076; (CXXXI) & 1.077; (CXXXI) & 1.078; (CXXXI) & 1.079; (CXXXI) & 1.080; (CXXXI) & 1.081; (CXXXI) & 1.082; (CXXXI) & 1.083; (CXXXI) & 1.084; (CXXXI) & 1.085; (CXXXI) & 1.086; (CXXXI) & 1.087; (CXXXII) & 1.059; (CXXXII) & 1.060; (CXXXII) & 1.061; (CXXXII) & 1.062; (CXXXII) & 1.063; (CXXXII) & 1.064; (CXXXII) & 1.065; (CXXXII) & 1.066; (CXXXII) & 1.067;

(R2) & 1.067; (R2) & 1.068; (R2) & 1.069; (R2) & 1.070; (R2) & 1.071; (R2) & 1.072; (R2) & 1.073; (R2) & 1.074; (R2) & 1.075; (R2) & 1.076; (R2) & 1.077; (R2) & 1.078; (R2) & 1.079; (R2) & 1.080; (R2) & 1.081; (R2) & 1.082; (R2) & 1.083; (R2) & 1.084; (R2) & 1.085; (R2) & 1.086; and (R2) & 1.087, preference being given to those combinations in which a partner is underlined.

The following reaction scheme gives a diagrammatic overview of the preparation of the compounds of formula (I):



The substituents R, R₁, R₂, R₂', X, Y, T and U indicated in the above scheme are as defined for formula (I); W is a leaving group; Hal is halogen, such as fluorine, chlorine, bromine or iodine, preferably chlorine, bromine or iodine.

Leaving groups W in the compounds of formulae (VI) and (VIII) that are suitable for the reactions are known from the literature and are described, for example, in: Houben-Weyl-Methoden der organischen Chemie, Vol. E4-carbonic acid derivatives (pages 149-165), H. Hagemann (Eds.), Georg Thieme Verlag, Stuttgart 1983. Especially preferred leaving groups are halogen, preferably iodine or chlorine; C₁-C₈alkoxy, C₁-C₈alkylthio, phenoxy, N-hydroxy-succinimide, N-hydroxy-phthalimide, imidazole, triazoles and 1-hydroxybenzotriazole-N-oxyl. It will be understood that all leaving groups that contain aliphatic or aromatic rings may be unsubstituted or substituted at those rings by customary substituents.

The compounds of formula (XX) are likewise novel and exhibit a pesticidal spectrum of action similar to that of the compounds of formula (I). The present invention relates to

those compounds also. In the compounds of formula (XX), as regards the substituents R_1 , R_2 , R_2' , X, Y, T and U preference is given to the same substituents as those already mentioned for the preferred sub-groups of compounds of formula (I), Hal preferably being fluorine or chlorine, especially chlorine. As a result of the Hal substituent, these novel compounds are excellently suitable for further derivatisation and therefore for the preparation of parasiticides, for example the compounds of formula (I).

The reactions illustrated and described hereinabove and hereinbelow are carried out in a manner known *per se*, for example in the absence or, usually, in the presence of a suitable solvent or diluent or a mixture thereof, the operation being carried out, as required, with cooling, at room temperature or with heating, for example in a temperature range of from about -80°C to the boiling temperature of the reaction medium, preferably from about -40°C to about $+120^{\circ}\text{C}$, especially from -20°C to 40°C and, if necessary, in a closed vessel, under pressure, in an inert gas atmosphere and/or under anhydrous conditions. Especially advantageous reaction conditions for each individual reaction step can be found in the explanations which follow and in the specific Preparation Examples.

Unless special conditions are mentioned, the reactants can in principle be reacted with one another as such, that is to say without the addition of a solvent or diluent, for example in the molten state. It is more advantageous, however, to add an inert solvent or diluent or a mixture thereof. Examples of such solvents or diluents that may be mentioned include: aromatic, aliphatic and alicyclic hydrocarbons and halogenated hydrocarbons, such as benzene, toluene, xylene, mesitylene, Tetralin, chlorobenzene, dichlorobenzene, bromobenzene, petroleum ether, hexane, cyclohexane, dichloromethane, trichloromethane, tetrachloromethane, dichloroethane, trichloroethene or tetrachloroethene; esters, such as ethyl acetate; ethers, such as diethyl ether, dipropyl ether, diisopropyl ether, dibutyl ether, tert-butyl methyl ether, ethylene glycol monomethyl ether, ethylene glycol monoethyl ether, ethylene glycol dimethyl ether, dimethoxydiethyl ether, tetrahydrofuran or dioxane; ketones, such as acetone, methyl ethyl ketone or methyl isobutyl ketone; alcohols, such as methanol, ethanol, propanol, isopropanol, butanol, ethylene glycol or glycerol; amides, such as N,N-dimethylformamide, N,N-diethylformamide, N,N-dimethylacetamide, N-methylpyrrolidone or hexamethylphosphoric acid triamide; nitriles, such as acetonitrile or propionitrile; and sulfoxides, such as dimethyl sulfoxide. If the reaction in question is carried out in the presence of a base, bases used in excess, such as triethylamine, pyridine, N-methylmorpholine or

N,N-diethylaniline, may also serve as solvent or diluent. If the reaction is carried out in the presence of an acid catalyst, it is also possible to employ as solvent or diluent acids used in excess, e.g. strong organic carboxylic acids, such as unsubstituted or substituted, for example halo-substituted, C₁-C₄alkanecarboxylic acids, e.g. formic acid, acetic acid or propionic acid. Suitable solvents for the reaction in question can be taken from the Examples given below.

Bases suitable for facilitating those reactions in which base catalysts may optionally be used are, for example, alkali metal or alkaline earth metal hydroxides, hydrides, alkylides, amides, alkanolates, acetates, carbonates, dialkylamides or alkylsilylamides; alkylamines, alkylenediamines, free or N-alkylated, saturated or unsaturated cycloalkylamines, basic heterocycles, ammonium hydroxides and also carbocyclic amines. Examples are butyllithium, sodium hydroxide, sodium hydride, sodium amide, sodium methanolate, sodium acetate, sodium carbonate, potassium tert-butanolate, potassium hydroxide, potassium carbonate, potassium hydride, lithium diisopropylamide, potassium bis(trimethylsilyl)amide, calcium hydride, triethylamine, diisopropylethylamine, triethylenediamine, cyclohexylamine, N-cyclohexyl-N,N-dimethylamine, N,N-diethylaniline, pyridine, 4-(N,N-dimethylamino)pyridine, quinuclidine, N-methylmorpholine, benzyltrimethylammonium hydroxide and also 1,5-diazabicyclo[5.4.0]undec-5-ene (DBU). For replacing chlorine by iodine in a compound of formula (VI), the base used is preferably silver carbonate and the reagent used sodium iodide.

The procedure in detail is as follows: the compounds of formula (I) are prepared by either (a) reacting a compound of formula (IX) in an aprotic, advantageously polar, solvent in the presence of a suitable base and at relatively low temperatures with a compound of formula (VIII) or preferably (b) reacting a compound of formula (XX) with an acid RCOOH and isolating the end product from the reaction mixture, the substituents R, R₁, R₂, R₂', X, Y, T and U being as defined for formula (I); W being a leaving group; and Hal being halogen, such as fluorine, chlorine, bromine or iodine, preferably chlorine, bromine or iodine. The reaction is advantageously carried out in an anhydrous medium and under an inert gas atmosphere, for example under nitrogen or argon. The said reactions take place within a period of minutes up to several hours.

A preferred embodiment comprises the following features: the compound of formula (IX) is first dissolved in an anhydrous, aprotic, polar solvent, and then at a relatively low temperature, for example at ambient temperature or lower, an equimolar amount of one of the suitable bases described above, for example sodium hydride, is added and the mixture is stirred for a little longer at the same temperature. An equimolar amount of the compound of formula (VIII) dissolved in the same solvent is then added in portions, e.g. by dropwise addition, and stirring of the reaction mixture is continued at the same low temperature for a little longer. Any excess base is then neutralised and the reaction mixture is stirred for a few minutes longer and finally concentrated. The residue is advantageously taken up in a polar solvent, such as ethyl acetate, and optionally washed with water, and the organic phase is separated off and dried over a drying agent, for example an alkali or alkaline earth metal sulfate or carbonate, preferably magnesium or sodium sulfate, concentrated and purified. A suitable purification step is chromatography, for example on silica gel (ethyl acetate : hexane / 1 :1). Compounds of formula (I) are generally obtained in the form of colourless to dark yellow oils, resins or in the form of solids. The said oils and resins crystallise after a few days or weeks when stored, for example, in a freezer at about from -18° to -25°C. In the context of the present invention, relatively low temperatures are to be understood as being temperatures around room temperature and below or a temperature range of from about +25°C to about -80°C, preferably from room temperature to about -20°C.

The compounds of formula (IX) are known *per se* from the literature or can be prepared analogously to the examples described in the literature. For example, compounds of formula (IX) wherein Het is pyridyl that is unsubstituted or mono- or poly-substituted by halogen are described, together with their preparation, in European published specification EP-0 302 833. Further compounds of formula (IX) are disclosed in the following patent references, for example in European published specifications Nos. 285 985; 302 389, 376 279; 471 372; 364 844; 493 369; 381 130; 529 680; 163 855; 375 907; 259 738; 386 565; 383 091 and 590 425; US Patents 5 063 236; 5 302 605 and 4 742 060; and also in DE-4 207 604; GB-2 228 003 and WO 93/24002.

The compounds of formula (VIII) wherein R, R₂ and R₂' are as defined for formula (I) and W is one of the leaving groups mentioned above can be prepared by introducing the radical RCOO- into a compound of formula (VI). For that purpose it is advantageous to prepare a suspension of a compound of formula (VI) wherein Hal is iodine and an organic acid

R-COOH, an example of which is benzoic acid, and silver carbonate in an aprotic solvent, e.g. toluene or xylene. The suspension is heated at from about 50°C to about 100°C for a few hours and then the reaction mixture is cooled to room temperature and insoluble constituents are filtered off. The filtrate is washed with water and/or aqueous sodium chloride solution and dried over a customary drying agent, such as magnesium or sodium sulfate. On concentration, the compound of formula (VIII) is obtained in the form of an oil or a crystalline solid. It is generally unnecessary to carry out further purification before use in the next reaction step.

The preparation of compounds of formula (VI) wherein R_2 and R_2' are as defined for formula (I), W is one of the leaving groups mentioned hereinabove and Hal is iodine is effected by replacing by iodine the chlorine atom in a compound of formula (VI) wherein Hal is chlorine. For this purpose a suspension of the compound of formula (VI) wherein Hal is chlorine, an equimolar amount of sodium iodide and sodium hydrogen carbonate in a polar solvent, such as acetone, is prepared and is stirred at slightly elevated temperature, about 40°C, for from 12 to 24 hours. The resulting precipitate is filtered off and washed with acetone. The filtrate is concentrated, and diethyl ether and water are added. The organic phase is separated off and washed with aqueous potassium sulfite solution, then washed with aqueous sodium chloride solution and dried over a customary drying agent, such as magnesium or sodium sulfate. On concentration, the compound of formula (VI) is obtained in the form of a colourless, crystalline product, which can be filtered off and freed of solvent residues, e.g. *in vacuo*. Further purification before use in the next reaction step is unnecessary.


Compounds of formula (VI) can be prepared by dissolving compounds of formula (VI) wherein R_2 and R_2' are as defined for formula (I) and W is chlorine in a halogenated solvent, such as dichloromethane, and at low temperature, advantageously about 0°C, adding a solution of a compound of the formula H-W in portions and then stirring the reaction mixture at the low temperature for about 1 to 3 hours. Water is then added and the organic phase is separated off, washed with 1N sodium hydroxide solution and then with aqueous sodium chloride solution and dried over a drying agent, e.g. magnesium sulfate. On concentration, the compound of formula (VI) is obtained in the form of a colourless, crystalline product, which can be filtered off and freed of solvent residues *in vacuo*.

As illustrated in the above reaction scheme, the novel compounds of formula (XX) are prepared by reacting a compound of formula (IX) in an aprotic, advantageously polar, solvent in the presence of a suitable base and at relatively low temperatures, with a compound of formula (VI), the substituents R_1 , R_2 , R_2' , X, Y, T and U in the formulae (XX), (IX) and (VI) being as defined for formula (I); W being a leaving group; and Hal being halogen, such as fluorine, chlorine, bromine or iodine and preferably chlorine or iodine. The compound of formula (IX) is first dissolved in an anhydrous, aprotic, polar solvent, and then at relatively low temperature, e.g. ambient temperature or lower, an equimolar amount of one of the suitable bases described above, e.g. sodium hydride, is added and the mixture is stirred for a little longer at the same temperature. An equimolar amount of the compound of formula (VI) dissolved in the same solvent is then added in portions, e.g. by dropwise addition, and stirring of the reaction mixture is continued at the same low temperature for a little longer. Any excess base is then neutralised and the reaction mixture is stirred for a few minutes longer and finally concentrated. The residue is advantageously taken up in a polar solvent, such as ethyl acetate, and optionally washed with water, and the organic phase is separated off and dried over a drying agent, for example an alkali or alkaline earth metal sulfate or carbonate, preferably magnesium or sodium sulfate, concentrated and purified. A suitable purification step is chromatography, for example on silica gel (ethyl acetate : hexane / 1 : 1). Compounds of formula (XX) are generally obtained in the form of colourless to dark yellow oils, resins or, predominantly, in the form of solids.

Preparation examples

Preparation of preliminary products


Preparation of carbonic acid (chloromethyl ester) (4-nitrophenyl ester):



At 0°C, a solution of 69.6 g of 4-nitrophenol and 39.6 g of pyridine in 500 ml of dichloromethane is added dropwise in the course of 30 minutes to a solution of 71 g of chloromethyl chloroformate in 1000 ml of dichloromethane. The reaction mixture is stirred at 0°C for a further 2 hours and then 1000 ml of H₂O are added. The organic phase is washed using 250 ml each of 1N NaOH and aqueous sodium chloride solution and dried over MgSO₄.


After concentration of the solvent, 111 g of a white solid having a melting point of 61-62°C are obtained.

Preparation of carbonic acid (iodomethyl ester) (4-nitrophenyl ester)




34.8 g of carbonic acid (chloromethyl ester) (4-nitrophenyl ester), 45.0 g of NaI and 2.52 g of NaHCO₃ are suspended in 350 ml of acetone and the resulting suspension is stirred at a temperature of 40°C for 16 hours. The precipitate is filtered off and washed with 100 ml of acetone. The filtrate is concentrated and the residue is taken up in 500 ml of diethyl ether and 100 ml of H₂O. The organic phase is washed with 100 ml each of saturated potassium disulfite solution and aqueous sodium chloride solution, dried over MgSO₄ and concentrated. 47.4 g of a white solid having a melting point of 45-46°C are obtained.

Preparation of benzoic acid 4-nitrophenoxycarbonyloxymethyl ester:



0.97 g of carbonic acid (iodomethyl ester) (4-nitrophenyl ester), 0.55 g of benzoic acid and 1.24 g of Ag₂CO₃ are suspended in 30 ml of toluene and stirred at 80°C for 3 hours. The mixture is cooled to room temperature and the precipitate is filtered off. The filtrate is washed with 15 ml each of H₂O and aqueous sodium chloride solution, dried over MgSO₄ and concentrated. 0.75 g of the desired product is obtained in the form of a white solid having a melting point of 69-71°C.


Preparation of carbonic acid (chloromethyl ester) (N-hydroxysuccinimide ester):



At 0°C, a solution of 99.3 g of N-hydroxysuccinimide and 55.4 g of pyridine in 250 ml of dichloromethane is added in the course of 60 minutes to a solution of 80.6 g of

chloromethyl chloroformate in 750 ml of dichloromethane. The reaction mixture is stirred at 0°C for a further 5 hours and then 1000 ml of H₂O are added. The organic phase is washed with 250 ml each of 1N NaOH and aqueous sodium chloride solution and dried over MgSO₄. After concentration of the solvent, 119 g of a white solid having a melting point of 103-105°C are obtained.


Preparation of {1-[(6-chloro-pyridin-3-ylmethyl)-ethyl-amino]-2-nitro-vinyl}-methyl-carbamic acid) chloromethyl ester:



6.35 g of sodium hydride are added in portions at 0°C under a N₂ atmosphere to a solution of 65 g of N-(6-chloro-pyridin-3-ylmethyl)-N-ethyl-N'-methyl]-2-nitro-vinylidenediamine in 300 ml of DMF. The mixture is stirred at 0°C for 1 hour and then, at -20°C, 50 g of carbonic acid (chloromethyl ester) (N-hydroxysuccinimide ester) dissolved in 200 ml of DMF are added dropwise thereto. After 1 hour at -20°C, 250 ml of saturated NH₄Cl solution and 300 ml of ethyl acetate are added. The organic phase is separated off and washed with 300 ml of 1N NaOH, 300 ml of water and 300 ml of saturated sodium chloride solution. After drying over MgSO₄, the solvent system is concentrated and the residue is stirred at reflux temperature with 300 ml of MeOH. After cooling to room temperature, 33.5 g of whitish-yellow crystals having a melting point of 142-143°C are obtained.

Preparation of end products


Preparation of benzoic acid ({1-[(6-chloro-pyridin-3-ylmethyl)-ethyl-amino]-2-nitro-vinyl}-methyl-carbamoyloxy)-methyl ester:



Under a N₂ atmosphere, 53 mg of NaH are added to a solution of 270 mg of N-(6-chloro-pyridin-3-ylmethyl)-N-ethyl-N'-methyl]-2-nitro-vinylidenediamine in 20 ml of DMF. The mixture is stirred at room temperature for 1 hour. Then, at -20°C, 635 mg of benzoic acid 4-nitro-phenoxy-carbonyloxymethyl ester dissolved in 5 ml of DMF are added dropwise. After

6 hours at -20°C, 1 ml of saturated NH₄Cl solution is added and the solvent system is concentrated. The residue is taken up in 50 ml of ethyl acetate and washed with 25 ml of H₂O. The organic phase is dried over MgSO₄ and concentrated, and the residue is chromatographed on silica gel (ethyl acetate:hexane; 1:1). 450 mg of whitish-yellow crystals having a melting point of 62-63°C are obtained.

Preparation of 2,2-dimethylbutyric acid {1-[(6-chloro-pyridin-3-ylmethyl)-ethyl-amino]-2-nitro-vinyl)-methyl-carbamoyloxy)-methyl ester:




A mixture of 10.9 g of {1-[(6-chloro-pyridin-3-ylmethyl)-ethyl-amino]-2-nitro-vinyl)-methyl-carbamoyl acid chloromethyl ester, 4.18 g of 2,2-dimethylbutyric acid and 5.39 g of potassium carbonate in 125 ml of DMF is stirred at a temperature of 50°C for 12 hours. The mixture is cooled to room temperature, the precipitate is filtered off and 250 ml of diethyl ether and 250 ml of 1N NaOH are added to the filtrate. The organic phase is separated off and washed with 250 ml of water and 250 ml of saturated sodium chloride solution. After drying over MgSO₄, the solvent system is concentrated and the residue is stirred with 200 ml of diethyl ether and 100 ml of hexane. 11.1 g of white crystals having a melting point of 79-80°C are obtained.

In an analogous manner it is also possible to prepare the following typical compounds listed in the Table.

Typical compounds of formula (I) are listed in Tables 1 to 4, while Tables 5 to 8 give typical examples of intermediates of formula (XX). The present invention is not limited to the typical examples shown, however.

Table 1: Compounds of formula (Xa)



No.	M	L	T	U	R ₂	R ₂ '	R	physical data
1.001	6-Cl	H	C ₂ H ₅	CH ₃	H	H	CH ₃	yellow resin
1.002	6-Cl	5-Cl	C ₂ H ₅	CH ₃	H	H	CH ₃	
1.003	6-Cl	H	CH ₃	CH ₃	H	H	CH ₃	
1.004	6-Cl	H	C ₂ H ₅	CH ₃	CH ₃	H	CH ₃	
1.005	6-Cl	H	C ₂ H ₅	CH ₃	H	H	C ₂ H ₅	
1.006	6-Cl	H	C ₂ H ₅	CH ₃	H	H	i-C ₃ H ₇	
1.007	6-Cl	H	C ₂ H ₅	CH ₃	H	H	n-C ₃ H ₇	
1.008	6-Cl	H	C ₂ H ₅	CH ₃	H	H	n-C ₄ H ₉	m.p. 66-68°C
1.009	6-Cl	H	H	CH ₃	H	H	s-C ₄ H ₉	
1.010	H	5-Cl	C ₂ H ₅	CH ₃	H	H	t-C ₄ H ₉	
1.011	6-Cl	H	C ₂ H ₅	CH ₃	H	H	n-C ₇ H ₁₅	m.p. 84-85°C
1.012	6-Cl	H	C ₂ H ₅	CH ₃	H	H	n-C ₁₁ H ₂₃	yellow resin
1.013	6-Cl	H	C ₂ H ₅	CH ₃	H	H	n-C ₁₃ H ₂₇	m.p. 52-54°C
1.014	6-Cl	H	C ₂ H ₅	CH ₃	H	H	n-C ₁₅ H ₃₁	m.p. 56-58°C
1.015	6-Cl	H	C ₂ H ₅	CH ₃	H	H	n-C ₁₇ H ₃₅	m.p. 60-61°C
1.016	6-Cl	5-Cl	C ₂ H ₅	CH ₃	H	H	n-C ₁₁ H ₂₃	
1.017	6-Cl	5-F	CH ₃	CH ₃	H	H	n-C ₇ H ₁₅	
1.018	6-Cl	H	C ₂ H ₅	CH ₃	H	H	phenyl	glassy m.p. 62-63°C
1.019	6-Cl	H	C ₂ H ₅	CH ₃	H	H	n-C ₅ H ₁₁	resin

1.020	6-Cl	H	C ₂ H ₅	CH ₃	H	H	biphenyl	glassy m.p. 72-69°C glassy m.p. 60-61°C glassy m.p. 103-5°C
1.021	6-Cl	H	C ₂ H ₅	CH ₃	H	H	phenoxyphenyl	
1.022	6-Cl	H	C ₂ H ₅	CH ₃	H	H	4-t-butyl-phenyl	
1.023	6-Cl	H	C ₂ H ₅	CH ₃	H	H	2-chloro-phenyl	
1.024	6-Cl	H	C ₂ H ₅	CH ₃	H	H	2,6-difluoro-phenyl	
1.025	6-Cl	H	C ₂ H ₅	H	H	H	n-C ₁₇ H ₃₅	
1.026	6-Cl	H	C ₂ H ₅	C ₂ H ₅	CH ₃	H	n-C ₇ H ₁₅	
1.027	6-Cl	H	C ₂ H ₅	CH ₃	C ₂ H ₅	H	CH ₃	
1.028	6-Cl	H	C ₂ H ₅	CH ₃	n-C ₃ H ₇	H	CH ₃	
1.029	6-Cl	H	C ₂ H ₅	CH ₃	s-C ₄ H ₉	H	CH ₃	
1.030	6-Cl	H	C ₂ H ₅	CH ₃	n-C ₆ H ₁₃	H	CH ₃	
1.031	6-Cl	H	CH ₃	CH ₃	CH ₃	H	CH ₃	
1.032	6-Cl	4-F	CH ₃	CH ₃	CH ₃	H	CH ₃	
1.033	6-Cl	H	C ₂ H ₅	CH ₃	H	H	cyclohexyl	
1.034	6-Cl	H	C ₂ H ₅	CH ₃	H	H	vinyl	
1.035	6-Cl	H	C ₂ H ₅	CH ₃	H	H	allyl	
1.036	6-Cl	H	C ₂ H ₅	CH ₃	H	H	2-propargyl	
1.037	6-Cl	H	C ₂ H ₅	CH ₃	CH ₃	CH ₃	CH ₃	
1.038	6-Cl	H	C ₂ H ₅	CH ₃	CH ₃	CH ₃	n-C ₄ H ₉	
1.039	6-Cl	H	C ₂ H ₅	CH ₃	CH ₃	CH ₃	n-C ₇ H ₁₅	
1.040	6-Cl	H	C ₂ H ₅	CH ₃	C ₂ H ₅	CH ₃	n-C ₁₁ H ₂₃	
1.041	6-Cl	H	C ₂ H ₅	CH ₃	CH ₃	CH ₃	n-C ₁₃ H ₂₇	
1.042	6-Cl	H	C ₂ H ₅	CH ₃	C ₂ H ₅	C ₂ H ₅	n-C ₁₅ H ₃₁	
1.043	6-Cl	H	C ₂ H ₅	CH ₃	n-C ₃ H ₇	C ₂ H ₅	n-C ₁₇ H ₃₅	
1.044	6-Cl	H	C ₂ H ₅	CH ₃	CH ₃	s-C ₄ H ₉	phenyl	

1.045	6-Cl	H	C ₂ H ₅	CH ₃	CH ₃	n-C ₃ H ₇	biphenyl	
1.046	6-Cl	H	C ₂ H ₅	CH ₃	s-C ₄ H ₉	C ₂ H ₅	phenoxyphenyl	
1.047	6-Cl	H	C ₂ H ₅	CH ₃	s-C ₄ H ₉	C ₂ H ₅	4-s-butyl-phenyl	
1.048	6-Cl	H	C ₂ H ₅	CH ₃	s-C ₄ H ₉	n-C ₃ H ₇	2-fluoro-phenyl	
1.049	6-Cl	H	C ₂ H ₅	C ₂ H ₅	CH ₃	n-C ₃ H ₇	n-C ₇ H ₁₅	
1.050	6-Cl	H	C ₂ H ₅	CH ₃	C ₂ H ₅	CH ₃	CH ₃	
1.051	6-Cl	H	C ₂ H ₅	CH ₃	n-C ₃ H ₇	CH ₃	CH ₃	
1.052	6-Cl	H	C ₂ H ₅	CH ₃	s-C ₄ H ₉	CH ₃	CH ₃	
1.053	6-Cl	H	C ₂ H ₅	CH ₃	n-C ₆ H ₁₃	H	CH ₃	
1.054	6-Cl	H	C ₂ H ₅	CH ₃	H	H	C(CH ₃) ₂ C ₃ H ₇ -n	m.p. 73-74°C
1.055	6-Cl	H	C ₂ H ₅	CH ₃	H	H	C(CH ₃) ₂ C ₃ H ₇ -n	m.p. 92-93°C
1.056	6-Cl	H	C ₂ H ₅	CH ₃	H	H	C(CH ₃) ₂ C ₂ H ₅	m.p. 79-80°C
1.057	6-Cl	H	C ₂ H ₅	CH ₃	H	H	n-C ₆ H ₁₃	m.p. 84-85°C
1.058	6-Cl	H	C ₂ H ₅	CH ₃	H	H	n-C ₁₀ H ₂₁	resin
1.059	6-Cl	H	C ₂ H ₅	CH ₃	H	H	CH(CH ₃)C ₃ H ₇ n	m.p. 73-74°C
1.060	6-Cl	H	C ₂ H ₅	CH ₃	H	H	n-C ₅ H ₁₁	oil
1.061	6-Cl	H	C ₂ H ₅	CH ₃	H	H	n-C ₈ H ₁₇	m.p. 53-55°C
1.062	6-Cl	H	C ₂ H ₅	CH ₃	H	H	n-C ₄ H ₉	oil
1.063	6-Cl	H	C ₂ H ₅	CH ₃	H	H	1-phenyl-1-cyclopentyl	m.p. 101-103°C
1.064	6-Cl	H	C ₂ H ₅	CH ₃	H	H	1-phenyl-1-cyclopropyl	glass
1.065	6-Cl	H	C ₂ H ₅	CH ₃	H	H	1-phenyl-1-cyclohexyl	m.p. 110-112°C
1.066	6-Cl	H	C ₂ H ₅	CH ₃	H	H	C(CH ₃) ₂ phenyl	m.p. 104-105°C
1.067	6-Cl	H	C ₂ H ₅	CH ₃	H	H	2,6-dimethyl-phenyl	m.p. 108-111°C
1.068	6-Cl	H	C ₂ H ₅	CH ₃	H	H	2,4,6-tri-isopropyl-phenyl	m.p. 118-119°C
1.069	6-Cl	H	C ₂ H ₅	CH ₃	H	H	cyclohexyl	m.p. 94-96°C

1.070	6-Cl	H	C ₂ H ₅	CH ₃	H	H	benzyl	glass
1.071	6-Cl	H	C ₂ H ₅	CH ₃	H	H	1-adamantyl	m.p. 158-160°C
1.072	6-Cl	H	C ₂ H ₅	CH ₃	H	H	t-C ₄ H ₉	m.p. 136-137°C
1.073	6-Cl	H	C ₂ H ₅	CH ₃	H	H	CH(phenyl) ₂	glass
1.074	6-Cl	H	C ₂ H ₅	CH ₃	H	H	cyclopropyl	m.p. 112-113°C
1.075	6-Cl	H	C ₂ H ₅	CH ₃	H	H	CH(C ₂ H ₅) ₂	m.p. 69-70°C
1.076	6-Cl	H	C ₂ H ₅	CH ₃	H	H	CH(n-C ₃ H ₇) ₂	m.p. 81-82°C
1.077	6-Cl	H	C ₂ H ₅	CH ₃	H	H	CH ₂ (cyclohexyl)	m.p. 71-73°C
1.078	6-Cl	H	C ₂ H ₅	CH ₃	H	H	CH ₂ CH ₂ (cyclohexyl)	m.p. 91-93°C
1.079	6-Cl	H	C ₂ H ₅	CH ₃	H	H	3-(t-C ₄ H ₉)cyclohex-1-yl	m.p. 107-110°C
1.080	6-Cl	H	C ₂ H ₅	CH ₃	H	H	1-(4-chlorophenyl)-1-cyclopentyl	m.p. 125-126°C
1.081	6-Cl	H	C ₂ H ₅	CH ₃	H	H	1-(4-fluorophenyl)-1-cyclopentyl	m.p. 105-107°C
1.082	6-Cl	H	C ₂ H ₅	CH ₃	H	H	CH ₂ C(CH ₃) ₂ CH ₃	m.p. 107-108°C
1.083	6-Cl	H	C ₂ H ₅	CH ₃	H	H	CH ₂ CH(phenyl) ₂	m.p. 91-93°C
1.084	6-Cl	H	C ₂ H ₅	CH ₃	H	H	1-methyl-2,2-dichloro-1-cyclopropyl	m.p. 80-82°C
1.085	6-Cl	H	C ₂ H ₅	CH ₃	H	H	1-methyl-1-cyclohexyl	m.p. 95-96°C
1.086	6-Cl	H	C ₂ H ₅	CH ₃	H	H	cyclopentyl	m.p. 134-135°C
1.087	6-Cl	H	C ₂ H ₅	CH ₃	CH ₃	CH ₃	4-Cl-phenyl	glass
1.088	6-Cl	H	CH ₃	CH ₃	CH ₃	CH ₃	CH ₃	
1.089	6-Cl	5-Cl	CH ₃	CH ₃	H	H	CH ₃	
1.090	6-Cl	H	CH ₃	CH ₃	H	H	CH ₃	
1.091	6-Cl	H	CH ₃	CH ₃	CH ₃	H	CH ₃	
1.092	6-Cl	H	CH ₃	CH ₃	H	H	C ₂ H ₅	
1.093	6-Cl	H	CH ₃	CH ₃	H	H	i-C ₃ H ₇	

1.094	6-Cl	H	CH ₃	CH ₃	H	H	n-C ₃ H ₇
1.095	6-Cl	H	CH ₃	CH ₃	H	H	n-C ₄ H ₉
1.096	6-Cl	H	CH ₃	CH ₃	H	H	s-C ₄ H ₉
1.097	H	5-Cl	CH ₃	CH ₃	H	H	t-C ₄ H ₉
1.098	6-Cl	H	CH ₃	CH ₃	H	H	n-C ₇ H ₁₅
1.099	6-Cl	H	CH ₃	CH ₃	H	H	n-C ₁₁ H ₂₃
1.100	6-Cl	H	CH ₃	CH ₃	H	H	n-C ₁₃ H ₂₇
1.101	6-Cl	H	CH ₃	CH ₃	H	H	n-C ₁₅ H ₃₁
1.102	6-Cl	H	CH ₃	CH ₃	H	H	n-C ₁₇ H ₃₅
1.103	6-Cl	5-Cl	CH ₃	CH ₃	H	H	n-C ₁₁ H ₂₃
1.104	6-Cl	5-F	CH ₃	CH ₃	CH ₃	CH ₃	n-C ₇ H ₁₅
1.105	6-Cl	H	CH ₃	CH ₃	H	H	phenyl
1.106	6-Cl	H	CH ₃	CH ₃	H	H	n-C ₅ H ₁₁
1.107	6-Cl	H	CH ₃	CH ₃	H	H	biphenyl
1.108	6-Cl	H	CH ₃	CH ₃	H	H	phenoxyphenyl
1.109	6-Cl	H	CH ₃	CH ₃	H	H	4-t-butyl-phenyl
1.110	6-Cl	H	CH ₃	CH ₃	H	H	2-chloro-phenyl
1.111	6-Cl	H	CH ₃	CH ₃	H	H	2,6-difluoro-phenyl
1.112	6-Cl	H	CH ₃	CH ₃	H	H	C(CH ₃) ₂ C ₃ H ₇ -n
1.113	6-Cl	H	CH ₃	CH ₃	H	H	C(CH ₃) ₂ C ₃ H ₇ -n
1.114	6-Cl	H	CH ₃	CH ₃	H	H	C(CH ₃) ₂ C ₂ H ₅
1.115	6-Cl	H	CH ₃	CH ₃	H	H	n-C ₆ H ₁₃
1.116	6-Cl	H	CH ₃	CH ₃	H	H	n-C ₁₀ H ₂₁
1.117	6-Cl	H	CH ₃	CH ₃	H	H	CH(CH ₃)C ₃ H ₇ -n
1.118	6-Cl	H	CH ₃	CH ₃	H	H	n-C ₅ H ₁₁


1.119	6-Cl	H	CH ₃	CH ₃	H	H	n-C ₈ H ₁₇
1.120	6-Cl	H	CH ₃	CH ₃	H	H	n-C ₄ H ₉
1.121	6-Cl	H	CH ₃	CH ₃	H	H	1-phenyl-1-cyclopentyl
1.122	6-Cl	H	CH ₃	CH ₃	H	H	1-phenyl-1-cyclopropyl
1.123	6-Cl	H	CH ₃	CH ₃	H	H	1-phenyl-1-cyclohexyl
1.124	6-Cl	H	CH ₃	CH ₃	H	H	C(CH ₃) ₂ phenyl
1.125	6-Cl	H	CH ₃	CH ₃	H	H	2,6-dimethylphenyl
1.126	6-Cl	H	CH ₃	CH ₃	H	H	2,4,6-tri-isopropylphenyl
1.127	6-Cl	H	CH ₃	CH ₃	H	H	cyclohexyl
1.128	6-Cl	H	CH ₃	CH ₃	H	H	benzyl
1.129	6-Cl	H	CH ₃	CH ₃	H	H	1-adamantyl
1.130	6-Cl	H	CH ₃	CH ₃	H	H	t-C ₄ H ₉
1.131	6-Cl	H	CH ₃	CH ₃	H	H	CH(phenyl) ₂
1.132	6-Cl	H	CH ₃	CH ₃	H	H	cyclopropyl
1.133	6-Cl	H	CH ₃	CH ₃	H	H	CH(C ₂ H ₅) ₂
1.134	6-Cl	H	CH ₃	CH ₃	H	H	CH(n-C ₃ H ₇) ₂
1.135	6-Cl	H	CH ₃	CH ₃	H	H	CH ₂ (cyclohexyl)
1.136	6-Cl	H	CH ₃	CH ₃	H	H	CH ₂ CH ₂ (cyclohexyl)
1.137	6-Cl	H	CH ₃	CH ₃	H	H	3-(t-C ₄ H ₉)cyclohex-1-yl
1.138	6-Cl	H	CH ₃	CH ₃	H	H	1-(4-chlorophenyl)-1-cyclopentyl
1.139	6-Cl	H	CH ₃	CH ₃	H	H	1-(4-fluorophenyl)-1-cyclopentyl
1.140	6-Cl	H	CH ₃	CH ₃	H	H	CH ₂ C(CH ₃) ₂ CH ₃
1.141	6-Cl	H	CH ₃	CH ₃	H	H	CH ₂ CH(phenyl) ₂
1.142	6-Cl	H	CH ₃	CH ₃	H	H	1-methyl-2,2-dichloro-1-cyclopropyl

1.143	6-Cl	H	CH ₃	CH ₃	H	H	1-methyl-1-cyclohexyl
1.144	6-Cl	H	CH ₃	CH ₃	H	H	cyclopentyl
1.145	6-Cl	H	CH ₃	CH ₃	CH ₃	CH ₃	4-Cl-phenyl
1.146	6-Cl	H	C ₂ H ₅	C ₂ H ₅	CH ₃	CH ₃	CH ₃
1.147	6-Cl	5-Cl	C ₂ H ₅	C ₂ H ₅	H	H	CH ₃
1.148	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	CH ₃
1.149	6-Cl	H	C ₂ H ₅	C ₂ H ₅	CH ₃	C ₂ H ₅	CH ₃
1.150	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	C ₂ H ₅
1.151	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	i-C ₃ H ₇
1.152	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₃ H ₇
1.153	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₄ H ₉
1.154	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	s-C ₄ H ₉
1.155	H	5-Cl	C ₂ H ₅	C ₂ H ₅	H	H	t-C ₄ H ₉
1.156	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₇ H ₁₅
1.157	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₁₁ H ₂₃
1.158	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₁₃ H ₂₇
1.159	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₁₅ H ₃₁
1.160	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₁₇ H ₃₅
1.161	6-Cl	5-Cl	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₁₁ H ₂₃
1.162	6-Cl	5-F	C ₂ H ₅	C ₂ H ₅	CH ₃	CH ₃	n-C ₇ H ₁₅
1.163	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	phenyl
1.164	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₅ H ₁₁
1.165	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	biphenyl
1.166	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	phenoxyphenyl
1.167	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	4-t-butyl-phenyl

1.168	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	2-chloro-phenyl
1.169	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	2,6-difluoro-phenyl
1.170	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	C(CH ₃) ₂ C ₃ H _{7-n}
1.171	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	C(CH ₃) ₂ C ₃ H _{7-n}
1.172	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	C(CH ₃) ₂ C ₂ H ₅
1.173	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₆ H ₁₃
1.174	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₁₀ H ₂₁
1.175	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	CH(CH ₃)C ₃ H _{7-n}
1.176	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₅ H ₁₁
1.177	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₈ H ₁₇
1.178	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₄ H ₉
1.179	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	1-phenyl-1-cyclopentyl
1.180	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	1-phenyl-1-cyclopropyl
1.191	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	1-phenyl-1-cyclohexyl
1.192	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	C(CH ₃) ₂ phenyl
1.193	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	2,6-dimethyl-phenyl
1.194	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	2,4,6-tri-isopropyl-phenyl
1.195	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	cyclohexyl
1.196	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	benzyl
1.197	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	1-adamantyl
1.198	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	t-C ₄ H ₉
1.199	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	CH(phenyl) ₂
1.200	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	cyclopropyl
1.201	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	CH(C ₂ H ₅) ₂
1.202	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	CH(n-C ₃ H ₇) ₂

1.203	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	CH ₂ (cyclohexyl)
1.204	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	CH ₂ CH ₂ (cyclohexyl)
1.205	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	3-(t-C ₄ H ₉)cyclohex-1-yl
1.206	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	1-(4-chlorophenyl)-1-cyclopentyl
1.207	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	1-(4-fluorophenyl)-1-cyclopentyl
1.208	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	CH ₂ C(CH ₃) ₂ CH ₃
1.209	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	CH ₂ CH(phenyl) ₂
1.210	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	1-methyl-2,2-dichloro-1-cyclopropyl
1.211	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	1-methyl-1-cyclohexyl
1.212	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	cyclopentyl
1.213	6-Cl	H	C ₂ H ₅	C ₂ H ₅	CH ₃	CH ₃	4-Cl-phenyl

Table 2: Compounds of formula (Xb)



No.	L	M	T	U	R ₂	R ₂ '	R
2.001	H	Cl	C ₂ H ₅	CH ₃	H	H	CH ₃
2.002	Cl	Cl	C ₂ H ₅	CH ₃	H	H	CH ₃
2.003	H	F	CH ₃	CH ₃	H	H	CH ₃
2.004	H	Cl	C ₂ H ₅	CH ₃	CH ₃	H	CH ₃
2.005	F	Cl	C ₂ H ₅	CH ₃	H	H	C ₂ H ₅
2.006	H	Cl	C ₂ H ₅	CH ₃	H	H	i-C ₃ H ₇
2.007	H	Cl	C ₂ H ₅	CH ₃	H	H	n-C ₃ H ₇

2.008	H	Cl	C ₂ H ₅	CH ₃	H	H	n-C ₄ H ₉
2.009	H	Cl	H	CH ₃	H	H	s-C ₄ H ₉
2.010	H	Cl	C ₂ H ₅	CH ₃	H	H	t-C ₄ H ₉
2.011	H	Cl	C ₂ H ₅	CH ₃	H	H	n-C ₇ H ₁₅
2.012	H	Cl	C ₂ H ₅	CH ₃	H	H	n-C ₁₁ H ₂₃
2.013	H	Cl	C ₂ H ₅	CH ₃	H	H	n-C ₁₃ H ₂₇
2.014	H	Cl	C ₂ H ₅	CH ₃	H	H	n-C ₁₅ H ₃₁
2.015	H	Cl	C ₂ H ₅	CH ₃	H	H	n-C ₁₇ H ₃₅
2.016	Cl	Cl	C ₂ H ₅	CH ₃	H	H	n-C ₁₁ H ₂₃
2.017	Cl	F	CH ₃	CH ₃	H	H	n-C ₇ H ₁₅
2.018	H	Cl	C ₂ H ₅	CH ₃	H	H	phenyl
2.019	H	Cl	C ₂ H ₅	CH ₃	H	CH ₃	phenyl
2.020	H	Cl	C ₂ H ₅	CH ₃	H	H	biphenyl
2.021	H	Cl	C ₂ H ₅	CH ₃	H	H	phenoxyphenyl
2.022	H	Cl	C ₂ H ₅	CH ₃	H	H	4-t-butyl-phenyl
2.023	H	Cl	C ₂ H ₅	CH ₃	H	H	2-chloro-phenyl
2.024	H	Cl	C ₂ H ₅	CH ₃	H	H	2,6-difluoro-phenyl
2.025	H	H	C ₂ H ₅	H	H	H	n-C ₁₇ H ₃₅
2.026	H	H	C ₂ H ₅	C ₂ H ₅	CH ₃	H	n-C ₇ H ₁₅
2.027	H	Cl	C ₂ H ₅	CH ₃	C ₂ H ₅	H	CH ₃
2.028	H	Cl	C ₂ H ₅	CH ₃	n-C ₃ H ₇	H	CH ₃
2.029	H	Cl	C ₂ H ₅	CH ₃	s-C ₄ H ₉	H	CH ₃
2.030	H	Cl	C ₂ H ₅	CH ₃	n-C ₆ H ₁₃	H	CH ₃
2.031	H	Cl	CH ₃	CH ₃	CH ₃	H	CH ₃
2.032	H	F	CH ₃	CH ₃	CH ₃	H	CH ₃

2.033	H	Cl	C ₂ H ₅	CH ₃	H	H	cyclohexyl
2.034	H	Cl	C ₂ H ₅	CH ₃	H	H	vinyl
2.035	H	Cl	C ₂ H ₅	CH ₃	H	H	allyl
2.036	H	Cl	C ₂ H ₅	CH ₃	H	H	2-propargyl
2.037	H	Cl	C ₂ H ₅	CH ₃	H	H	C(CH ₃) ₂ C ₃ H _{7-n}
2.038	H	Cl	C ₂ H ₅	CH ₃	H	H	C(CH ₃) ₂ C ₃ H _{7-n}
2.039	H	Cl	C ₂ H ₅	CH ₃	H	H	C(CH ₃) ₂ C ₂ H ₅
2.040	H	Cl	C ₂ H ₅	CH ₃	H	H	n-C ₆ H ₁₃
2.041	H	Cl	C ₂ H ₅	CH ₃	H	H	n-C ₁₀ H ₂₁
2.042	H	Cl	C ₂ H ₅	CH ₃	H	H	CH(CH ₃)C ₃ H _{7-n}
2.043	H	Cl	C ₂ H ₅	CH ₃	H	H	n-C ₅ H ₁₁
2.044	H	Cl	C ₂ H ₅	CH ₃	H	H	n-C ₈ H ₁₇
2.045	H	Cl	C ₂ H ₅	CH ₃	H	H	n-C ₄ H ₉
2.046	H	Cl	C ₂ H ₅	CH ₃	H	H	1-phenyl-1-cyclopentyl
2.047	H	Cl	C ₂ H ₅	CH ₃	H	H	1-phenyl-1-cyclopropyl
2.048	H	Cl	C ₂ H ₅	CH ₃	H	H	1-phenyl-1-cyclohexyl
2.049	2.0H	Cl	C ₂ H ₅	CH ₃	H	H	C(CH ₃) ₂ phenyl
2.050	H	Cl	C ₂ H ₅	CH ₃	H	H	2,6-dimethylphenyl
2.051	H	Cl	C ₂ H ₅	CH ₃	H	H	2,4,6-tri-isopropylphenyl
2.052	H	Cl	C ₂ H ₅	CH ₃	H	H	cyclohexyl
2.053	H	Cl	C ₂ H ₅	CH ₃	H	H	benzyl
2.054	H	Cl	C ₂ H ₅	CH ₃	H	H	1-adamantyl
2.055	H	Cl	C ₂ H ₅	CH ₃	H	H	t-C ₄ H ₉
2.056	H	Cl	C ₂ H ₅	CH ₃	H	H	CH(phenyl) ₂
2.057	H	Cl	C ₂ H ₅	CH ₃	H	H	cyclopropyl

2.058	H	Cl	C ₂ H ₅	CH ₃	H	H	CH(C ₂ H ₅) ₂
2.059	H	Cl	C ₂ H ₅	CH ₃	H	H	CH(n-C ₃ H ₇) ₂
2.060	H	Cl	C ₂ H ₅	CH ₃	H	H	CH ₂ (cyclohexyl)
2.061	H	Cl	C ₂ H ₅	CH ₃	H	H	CH ₂ CH ₂ (cyclohexyl)
2.062	H	Cl	C ₂ H ₅	CH ₃	H	H	3-(t-C ₄ H ₉)cyclohex-1-yl
2.063	H	Cl	C ₂ H ₅	CH ₃	H	H	1-(4-chloro-phenyl)-1-cyclopentyl
2.064	H	Cl	C ₂ H ₅	CH ₃	H	H	1-(4-fluoro-phenyl)-1-cyclopentyl
2.065	H	Cl	C ₂ H ₅	CH ₃	H	H	CH ₂ C(CH ₃) ₂ CH ₃
2.066	H	Cl	C ₂ H ₅	CH ₃	H	H	CH ₂ CH(phenyl) ₂
2.067	H	Cl	C ₂ H ₅	CH ₃	H	H	1-methyl-2,2-di-chloro-1-cyclopropyl
2.068	H	Cl	C ₂ H ₅	CH ₃	H	H	1-methyl-1-cyclohexyl
2.069	H	Cl	C ₂ H ₅	CH ₃	H	H	cyclopentyl
2.070	H	Cl	C ₂ H ₅	CH ₃	CH ₃	CH ₃	4-Cl-phenyl
2.071	6-Cl	H	CH ₃	CH ₃	CH ₃	CH ₃	CH ₃
2.073	6-Cl	5-Cl	CH ₃	CH ₃	H	H	CH ₃
2.074	6-Cl	H	CH ₃	CH ₃	H	H	CH ₃
2.075	6-Cl	H	CH ₃	CH ₃	CH ₃	H	CH ₃
2.076	6-Cl	H	CH ₃	CH ₃	H	H	C ₂ H ₅
2.077	6-Cl	H	CH ₃	CH ₃	H	H	i-C ₃ H ₇
2.078	6-Cl	H	CH ₃	CH ₃	H	H	n-C ₃ H ₇
2.079	6-Cl	H	CH ₃	CH ₃	H	H	n-C ₄ H ₉
2.080	6-Cl	H	CH ₃	CH ₃	H	H	s-C ₄ H ₉
2.081	H	5-Cl	CH ₃	CH ₃	H	H	t-C ₄ H ₉
2.082	6-Cl	H	CH ₃	CH ₃	H	H	n-C ₇ H ₁₅
2.083	6-Cl	H	CH ₃	CH ₃	H	H	n-C ₁₁ H ₂₃

2.084	6-Cl	H	CH ₃	CH ₃	H	H	n-C ₁₃ H ₂₇
2.085	6-Cl	H	CH ₃	CH ₃	H	H	n-C ₁₅ H ₃₁
2.086	6-Cl	H	CH ₃	CH ₃	H	H	n-C ₁₇ H ₃₅
2.087	6-Cl	5- Cl	CH ₃	CH ₃	H	H	n-C ₁₁ H ₂₃
2.088	6-Cl	5- F	CH ₃	CH ₃	CH ₃	CH ₃	n-C ₇ H ₁₅
2.089	6-Cl	H	CH ₃	CH ₃	H	H	phenyl
2.090	6-Cl	H	CH ₃	CH ₃	H	H	n-C ₅ H ₁₁
2.091	6-Cl	H	CH ₃	CH ₃	H	H	biphenyl
2.092	6-Cl	H	CH ₃	CH ₃	H	H	phenoxyphenyl
2.093	6-Cl	H	CH ₃	CH ₃	H	H	4-t-butyl-phenyl
2.094	6-Cl	H	CH ₃	CH ₃	H	H	2-chloro-phenyl
2.095	6-Cl	H	CH ₃	CH ₃	H	H	2,6-difluoro-phenyl
2.096	6-Cl	H	CH ₃	CH ₃	H	H	C(CH ₃) ₂ C ₃ H ₇ -n
2.097	6-Cl	H	CH ₃	CH ₃	H	H	C(CH ₃) ₂ C ₃ H ₇ -n
2.098	6-Cl	H	CH ₃	CH ₃	H	H	C(CH ₃) ₂ C ₂ H ₅
2.099	6-Cl	H	CH ₃	CH ₃	H	H	n-C ₆ H ₁₃
2.100	6-Cl	H	CH ₃	CH ₃	H	H	n-C ₁₀ H ₂₁
2.101	6-Cl	H	CH ₃	CH ₃	H	H	CH(CH ₃)C ₃ H ₇ n
2.102	6-Cl	H	CH ₃	CH ₃	H	H	n-C ₅ H ₁₁
2.103	6-Cl	H	CH ₃	CH ₃	H	H	n-C ₈ H ₁₇
2.104	6-Cl	H	CH ₃	CH ₃	H	H	n-C ₄ H ₉
2.105	6-Cl	H	CH ₃	CH ₃	H	H	1-phenyl-1-cyclo- pentyl
2.106	6-Cl	H	CH ₃	CH ₃	H	H	1-phenyl-1-cyclo- propyl
2.107	6-Cl	H	CH ₃	CH ₃	H	H	1-phenyl-1-cyclo- hexyl
2.108	6-Cl	H	CH ₃	CH ₃	H	H	C(CH ₃) ₂ phenyl


2.109	6-Cl	H	CH ₃	CH ₃	H	H	2,6-dimethylphenyl
2.110	6-Cl	H	CH ₃	CH ₃	H	H	2,4,6-tri-isopropyl-phenyl
2.111	6-Cl	H	CH ₃	CH ₃	H	H	cyclohexyl
2.112	6-Cl	H	CH ₃	CH ₃	H	H	benzyl
2.113	6-Cl	H	CH ₃	CH ₃	H	H	1-adamantyl
2.114	6-Cl	H	CH ₃	CH ₃	H	H	t-C ₄ H ₉
2.115	6-Cl	H	CH ₃	CH ₃	H	H	CH(phenyl) ₂
2.116	6-Cl	H	CH ₃	CH ₃	H	H	cyclopropyl
2.117	6-Cl	H	CH ₃	CH ₃	H	H	CH(C ₂ H ₅) ₂
2.118	6-Cl	H	CH ₃	CH ₃	H	H	CH(n-C ₃ H ₇) ₂
2.119	6-Cl	H	CH ₃	CH ₃	H	H	CH ₂ (cyclohexyl)
2.120	6-Cl	H	CH ₃	CH ₃	H	H	CH ₂ CH ₂ (cyclohexyl)
2.121	6-Cl	H	CH ₃	CH ₃	H	H	3-(t-C ₄ H ₉)cyclohex-1-yl
2.122	6-Cl	H	CH ₃	CH ₃	H	H	1-(4-chloro-phenyl)-1-cyclopentyl
2.123	6-Cl	H	CH ₃	CH ₃	H	H	1-(4-fluoro-phenyl)-1-cyclopentyl
2.124	6-Cl	H	CH ₃	CH ₃	H	H	CH ₂ C(CH ₃) ₂ CH ₃
2.125	6-Cl	H	CH ₃	CH ₃	H	H	CH ₂ CH(phenyl) ₂
2.126	6-Cl	H	CH ₃	CH ₃	H	H	1-methyl-2,2-di-chloro-1-cyclopropyl
2.127	6-Cl	H	CH ₃	CH ₃	H	H	1-methyl-1-cyclohexyl
2.128	6-Cl	H	CH ₃	CH ₃	H	H	cyclopentyl
2.129	6-Cl	H	CH ₃	CH ₃	CH ₃	CH ₃	4-Cl-phenyl
2.130	6-Cl	H	C ₂ H ₅	C ₂ H ₅	CH ₃	CH ₃	CH ₃
2.131	6-Cl	5-Cl	C ₂ H ₅	C ₂ H ₅	H	H	CH ₃
2.132	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	CH ₃
2.133	6-Cl	H	C ₂ H ₅	C ₂ H ₅	CH ₃	C ₂ H ₅	CH ₃

2.134	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	C ₂ H ₅
2.135	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	i-C ₃ H ₇
2.136	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₃ H ₇
2.137	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₄ H ₉
2.138	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	s-C ₄ H ₉
2.139	H	5- Cl	C ₂ H ₅	C ₂ H ₅	H	H	t-C ₄ H ₉
2.140	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₇ H ₁₅
2.141	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₁₁ H ₂₃
2.142	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₁₃ H ₂₇
2.143	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₁₅ H ₃₁
2.144	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₁₇ H ₃₅
2.145	6-Cl	5- Cl	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₁₁ H ₂₃
2.146	6-Cl	5- F	C ₂ H ₅	C ₂ H ₅	CH ₃	CH ₃	n-C ₇ H ₁₅
2.147	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	phenyl
2.148	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₅ H ₁₁
2.149	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	biphenyl
2.150	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	phenoxyphenyl
2.151	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	4-t-butyl-phenyl
2.152	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	2-chloro-phenyl
2.153	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	2,6-difluoro-phenyl
2.154	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	C(CH ₃) ₂ C ₃ H ₇ -n
2.155	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	C(CH ₃) ₂ C ₃ H ₇ -n
2.156	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	C(CH ₃) ₂ C ₂ H ₅
2.157	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₆ H ₁₃
2.158	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₁₀ H ₂₁

2.159	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	CH(CH ₃)C ₃ H ₇ n
2.160	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₅ H ₁₁
2.161	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₈ H ₁₇
2.162	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	n-C ₄ H ₉
2.163	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	1-phenyl-1-cyclo- pentyl
2.164	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	1-phenyl-1-cyclo- propyl
2.165	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	1-phenyl-1-cyclo- hexyl
2.166	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	C(CH ₃) ₂ phenyl
2.167	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	2,6-dimethylphenyl
2.168	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	2,4,6-tri-isopropyl- phenyl
2.169	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	cyclohexyl
2.170	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	benzyl
2.171	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	1-adamantyl
2.172	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	t-C ₄ H ₉
2.173	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	CH(phenyl) ₂
2.174	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	cyclopropyl
2.175	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	CH(C ₂ H ₅) ₂
2.176	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	CH(n-C ₃ H ₇) ₂
2.177	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	CH ₂ (cyclohexyl)
2.178	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	CH ₂ CH ₂ (cyclo- hexyl)
2.179	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	3-(t-C ₄ H ₉)cyclo- hex-1-yl
2.180	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	1-(4-chloro-phenyl)-1- cyclopentyl
2.181	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	1-(4-fluoro-phenyl)-1- cyclopentyl
2.182	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	CH ₂ C(CH ₃) ₂ CH ₃
2.183	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	CH ₂ CH(phenyl) ₂

2.184	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	1-methyl-2,2-di-chloro-1-cyclopropyl
2.185	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	1-methyl-1-cyclohexyl
2.186	6-Cl	H	C ₂ H ₅	C ₂ H ₅	H	H	cyclopentyl
2.187	6-Cl	H	C ₂ H ₅	C ₂ H ₅	CH ₃	CH ₃	4-Cl-phenyl

Table 3: Compounds of formula (XIa)




No.	Hal	Y	X	Q	R ₂	R ₂ '	R
3.001	Cl	NO ₂	CH	H	H	H	CH ₃
3.002	Cl	NO ₂	N	H	H	H	CH ₃
3.003	F	NO ₂	CH	H	H	H	CH ₃
3.004	Cl	NO ₂	CH	3-CH ₃	H	H	CH ₃
3.005	Cl	CN	CH	H	H	H	CH ₃
3.006	Cl	CN	N	H	H	H	CH ₃
3.007	Cl	NO ₂	CH	H	H	H	C ₂ H ₅
3.008	Cl	NO ₂	N	H	H	H	C ₂ H ₅
3.009	F	NO ₂	CH	H	H	H	C ₂ H ₅
3.010	Cl	NO ₂	CH	2-CH ₃	H	H	C ₂ H ₅
3.011	Cl	CN	CH	H	H	H	C ₂ H ₅
3.012	Cl	NO ₂	CH	H	H	H	n-C ₃ H ₇
3.013	Cl	NO ₂	N	H	H	H	n-C ₄ H ₉
3.014	F	NO ₂	CH	H	H	H	s-C ₄ H ₉
3.015	Cl	NO ₂	CH	3-CH ₃	H	H	t-C ₄ H ₉

3.016	Cl	CN	CH	H	H	H	n-C ₇ H ₁₅
3.017	Cl	CN	N	H	H	H	n-C ₁₁ H ₂₃
3.018	Cl	NO ₂	CH	H	H	H	n-C ₁₃ H ₂₇
3.019	Cl	NO ₂	N	H	H	H	n-C ₁₅ H ₃₁
3.020	F	NO ₂	CH	H	H	H	n-C ₁₇ H ₃₅
3.021	Cl	NO ₂	CH	2-CH ₃	H	H	n-C ₁₁ H ₂₃
3.022	Cl	CN	CH	H	H	H	n-C ₇ H ₁₅
3.023	Cl	NO ₂	CH	H	H	H	phenyl
3.024	Cl	NO ₂	N	H	H	CH ₃	phenyl
3.025	F	NO ₂	CH	H	H	H	biphenyl
3.026	Cl	NO ₂	CH	3-CH ₃	H	H	phenoxyphenyl
3.027	Cl	CN	CH	H	H	H	4-t-butyl-phenyl
3.028	Cl	CN	N	H	H	H	2-chloro-phenyl
3.029	Cl	NO ₂	CH	3-C ₂ H ₅	H	H	2,6-difluoro-phenyl
3.030	Cl	CN	CH	2-CH ₃	H	H	n-C ₁₇ H ₃₅
3.031	Cl	CN	N	3-CH ₃	CH ₃	H	n-C ₇ H ₁₅
3.032	Cl	CN	N	H	H	H	cyclohexyl
3.033	Cl	CN	N	H	H	H	vinyl
3.034	Cl	CN	CH	H	H	H	allyl
3.035	Cl	CN	N	H	H	H	2-propargyl
3.036	Cl	NO ₂	N	H	H	H	cyclohexyl
3.037	Cl	NO ₂	N	H	H	H	vinyl
3.038	Cl	NO ₂	N	H	H	H	allyl
3.039	Cl	NO ₂	CH	H	H	H	2-propargyl
3.040	Cl	CN	CH	H	C ₂ H ₅	H	C ₂ H ₅

3.041	Br	NO ₂	CH	H	H	H	n-C ₃ H ₇
3.042	Cl	NO ₂	N	H	C ₂ H ₅	CH ₃	n-C ₄ H ₉
3.043	F	NO ₂	CH	H	C ₂ H ₅	H	s-C ₄ H ₉
3.044	Br	NO ₂	CH	3-CH ₃	C ₂ H ₅	H	t-C ₄ H ₉
3.045	Cl	CN	CH	H	C ₂ H ₅	H	n-C ₇ H ₁₅
3.046	Cl	NO ₂	CH	H	H	H	C(CH ₃) ₂ C ₃ H ₇ -n
3.047	Cl	CN	CH	H	H	H	C(CH ₃) ₂ C ₃ H ₇ -n
3.048	Cl	NO ₂	CH	H	H	H	C(CH ₃) ₂ C ₂ H ₅
3.049	Cl	NO ₂	CH	H	H	H	n-C ₆ H ₁₃
3.050	Cl	NO ₂	CH	H	H	H	n-C ₁₀ H ₂₁
3.051	Cl	CN	C ₂ H ₅	CH ₃	H	H	CH(CH ₃)C ₃ H ₇ n
3.052	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	n-C ₅ H ₁₁
3.053	Cl	CN	C ₂ H ₅	CH ₃	H	H	n-C ₈ H ₁₇
3.054	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	n-C ₄ H ₉
3.055	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	1-phenyl-1-cyclopentyl
3.056	Cl	CN	C ₂ H ₅	CH ₃	H	H	1-phenyl-1-cyclopropyl
3.057	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	1-phenyl-1-cyclohexyl
3.058	Cl	CN	C ₂ H ₅	CH ₃	H	H	C(CH ₃) ₂ phenyl
3.059	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	2,6-dimethylphenyl
3.060	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	2,4,6-tri-isopropylphenyl
3.061	Cl	CN	C ₂ H ₅	CH ₃	H	H	cyclohexyl
3.062	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	benzyl
3.063	Cl	CN	C ₂ H ₅	CH ₃	H	H	1-adamantyl
3.064	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	t-C ₄ H ₉
3.065	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	CH(phenyl) ₂

3.066	Cl	CN	C ₂ H ₅	CH ₃	H	H	cyclopropyl
3.067	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	CH(C ₂ H ₅) ₂
3.068	Cl	CN	C ₂ H ₅	CH ₃	H	H	CH(n-C ₃ H ₇) ₂
3.069	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	CH ₂ (cyclohexyl)
3.070	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	CH ₂ CH ₂ (cyclohexyl)
3.071	Cl	CN	C ₂ H ₅	CH ₃	H	H	3-(1-C ₄ H ₉)cyclohex-1-yl
3.072	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	1-(4-chloro-phenyl)-1-cyclopentyl
3.073	Cl	CN	C ₂ H ₅	CH ₃	H	H	1-(4-fluoro-phenyl)-1-cyclopentyl
3.074	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	CH ₂ C(CH ₃) ₂ CH ₃
3.075	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	CH ₂ CH(phenyl) ₂
3.076	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	1-methyl-2,2-di-chloro-1-cyclopropyl
3.077	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	1-methyl-1-cyclohexyl
3.078	Cl	CN	C ₂ H ₅	CH ₃	H	H	cyclopentyl
3.079	Cl	NO ₂	C ₂ H ₅	CH ₃	CH ₃	CH ₃	4-Cl-phenyl

Table 4: Compounds of formula (XIIa)



No.	Hal	Y	X	T-U	R ₂	R ₂ '	R
4.001	Cl	NO ₂	CH	-(CH ₂) ₃ -	H	H	CH ₃
4.002	Cl	NO ₂	N	-(CH ₂) ₃ -	H	H	CH ₃
4.003	F	NO ₂	CH	-(CH ₂) ₃ -	H	H	CH ₃
4.004	F	NO ₂	N	-(CH ₂) ₃ -	H	H	CH ₃

4.005	Cl	CN	CH	-(CH ₂) ₃ -	H	H	CH ₃
4.006	Cl	CN	N	-(CH ₂) ₃ -	H	H	CH ₃
4.007	Cl	NO ₂	CH	-(CH ₂) ₃ -	H	H	C ₂ H ₅
4.008	Cl	NO ₂	N	-(CH ₂) ₃ -	H	H	C ₂ H ₅
4.009	F	NO ₂	CH	-(CH ₂) ₃ -	H	H	C ₂ H ₅
4.010	Cl	NO ₂	CH	-(CH ₂) ₃ -	CH ₃	H	C ₂ H ₅
4.011	Cl	CN	CH	-(CH ₂) ₃ -	H	H	C ₂ H ₅
4.012	Cl	NO ₂	CH	-(CH ₂) ₃ -	H	H	n-C ₃ H ₇
4.013	Cl	NO ₂	N	-(CH ₂) ₃ -	H	H	n-C ₄ H ₉
4.014	F	NO ₂	CH	-(CH ₂) ₃ -	H	H	s-C ₄ H ₉
4.015	Cl	NO ₂	CH	-(CH ₂) ₃ -	H	H	t-C ₄ H ₉
4.016	Cl	CN	CH	-(CH ₂) ₃ -	H	H	n-C ₇ H ₁₅
4.017	Cl	CN	N	-(CH ₂) ₃ -	H	H	n-C ₁₁ H ₂₃
4.018	Cl	NO ₂	CH	-(CH ₂) ₃ -	H	H	n-C ₁₃ H ₂₇
4.019	Cl	NO ₂	N	-(CH ₂) ₃ -	H	H	n-C ₁₅ H ₃₁
4.020	F	NO ₂	CH	-(CH ₂) ₃ -	H	H	n-C ₁₇ H ₃₅

4.021	Cl	NO ₂	CH	-(CH ₂) ₃ -	CH ₃	CH ₃	n-C ₁₁ H ₂₃
4.022	Cl	CN	CH	-(CH ₂) ₃ -	CH ₃	H	n-C ₇ H ₁₅
4.023	Cl	NO ₂	CH	-(CH ₂) ₃ -	H	H	phenyl
4.024	Cl	NO ₂	N	-(CH ₂) ₃ -	H	CH ₃	phenyl
4.025	F	CN	CH	-(CH ₂) ₃ -	H	H	biphenyl
4.026	Cl	NO ₂	CH	-(CH ₂) ₃ -	H	H	phenoxyphenyl
4.027	Cl	CN	CH	-(CH ₂) ₃ -	H	H	4-t-butyl-phenyl
4.028	Cl	CN	N	-(CH ₂) ₃ -	H	H	2-chloro-phenyl
4.029	Cl	NO ₂	CH	-(CH ₂) ₃ -	H	H	2,6-difluoro-phenyl
4.030	Cl	CN	CH	-(CH ₂) ₃ -	H	H	n-C ₁₇ H ₃₅
4.031	Br	CN	N	-(CH ₂) ₃ -	CH ₃	H	n-C ₇ H ₁₅
4.032	Cl	CN	N	-(CH ₂) ₃ -	H	H	cyclohexyl
4.033	Cl	CN	N	-(CH ₂) ₃ -	H	H	vinyl
4.034	Cl	CN	CH	-(CH ₂) ₃ -	H	H	allyl
4.035	Cl	CN	N	-(CH ₂) ₃ -	H	H	2-propargyl
4.036	Cl	NO ₂	N	-(CH ₂) ₃ -	H	H	cyclohexyl
4.037	Cl	NO ₂	N	-(CH ₂) ₃ -	H	H	vinyl

4.038	Cl	NO ₂	N	-(CH ₂) ₃ -	H	H	allyl
4.039	Cl	NO ₂	CH	-CH ₂ OCH ₂ -	H	H	2-propargyl
4.040	Cl	NO ₂	CH	-CH ₂ OCH ₂ -	H	H	CH ₃
4.041	Cl	NO ₂	N	-CH ₂ OCH ₂ -	H	H	CH ₃
4.042	F	NO ₂	CH	-CH ₂ OCH ₂ -	H	H	CH ₃
4.043	F	NO ₂	N	-CH ₂ OCH ₂ -	H	H	CH ₃
4.044	Cl	CN	CH	-CH ₂ OCH ₂ -	H	H	CH ₃
4.045	Cl	CN	N	-CH ₂ OCH ₂ -	H	H	CH ₃
4.045	Cl	NO ₂	CH	-CH ₂ OCH ₂ -	H	H	C ₂ H ₅
4.047	Cl	NO ₂	N	-CH ₂ OCH ₂ -	H	H	C ₂ H ₅
4.048	F	NO ₂	CH	-CH ₂ OCH ₂ -	H	H	C ₂ H ₅
4.049	Cl	NO ₂	CH	-CH ₂ OCH ₂ -	CH ₃	H	C ₂ H ₅
4.050	Cl	CN	CH	-CH ₂ OCH ₂ -	H	H	C ₂ H ₅
4.051	Cl	NO ₂	CH	-CH ₂ OCH ₂ -	H	H	n-C ₃ H ₇
4.052	Cl	NO ₂	N	-CH ₂ OCH ₂ -	H	H	n-C ₄ H ₉
4.053	F	NO ₂	CH	-CH ₂ OCH ₂ -	H	H	s-C ₄ H ₉


4.054	Cl	NO ₂	CH	-CH ₂ OCH ₂ -	H	H	t-C ₄ H ₉
4.055	Cl	CN	CH	-CH ₂ OCH ₂ -	H	H	n-C ₇ H ₁₅
4.056	Cl	CN	N	-CH ₂ OCH ₂ -	H	H	n-C ₁₁ H ₂₃
4.057	Cl	NO ₂	CH	-CH ₂ OCH ₂ -	H	H	n-C ₁₃ H ₂₇
4.058	Cl	NO ₂	N	-CH ₂ OCH ₂ -	H	H	n-C ₁₅ H ₃₁
4.059	F	NO ₂	CH	-CH ₂ OCH ₂ -	H	H	n-C ₁₇ H ₃₅
4.060	Cl	NO ₂	CH	-CH ₂ OCH ₂ -	CH ₃	CH ₃	n-C ₁₁ H ₂₃
4.061	Cl	CN	CH	-CH ₂ OCH ₂ -	CH ₃	H	n-C ₇ H ₁₅
4.062	Cl	NO ₂	CH	-CH ₂ OCH ₂ -	H	H	phenyl
4.063	Cl	NO ₂	N	-CH ₂ OCH ₂ -	H	CH ₃	phenyl
4.064	F	CN	CH	-CH ₂ OCH ₂ -	H	H	biphenyl
4.065	Cl	NO ₂	CH	-CH ₂ OCH ₂ -	H	H	phenoxyphenyl
4.066	Cl	CN	CH	-CH ₂ OCH ₂ -	H	H	4-t-butyl-phenyl
4.067	Cl	CN	N	-CH ₂ OCH ₂ -	H	H	2-chloro-phenyl
4.068	Cl	NO ₂	CH	-CH ₂ OCH ₂ -	H	H	2,6-difluoro-phenyl
4.069	Cl	CN	CH	-CH ₂ OCH ₂ -	H	H	n-C ₁₇ H ₃₅
4.070	Br	CN	N	-CH ₂ OCH ₂ -	CH ₃	H	n-C ₇ H ₁₅

4.071	Cl	CN	N	-CH ₂ OCH ₂ -	H	H	cyclohexyl
4.072	Cl	CN	N	-CH ₂ OCH ₂ -	H	H	vinyl
4.073	Cl	CN	CH	-CH ₂ OCH ₂ -	H	H	allyl
4.074	Cl	CN	N	-CH ₂ OCH ₂ -	H	H	2-propargyl
4.075	Cl	NO ₂	N	-CH ₂ OCH ₂ -	H	H	cyclohexyl
4.076	Cl	NO ₂	N	-CH ₂ OCH ₂ -	H	H	vinyl
4.077	Cl	NO ₂	N	-CH ₂ OCH ₂ -	H	H	allyl
4.078	Cl	NO ₂	CH	-CH ₂ OCH ₂ -	H	H	2-propargyl
4.079	Cl	NO ₂	CH	-CH ₂ N(CH ₃)CH ₂ -	H	H	CH ₃
4.080	Cl	NO ₂	N	-CH ₂ N(CH ₃)CH ₂ -	H	H	CH ₃
4.081	F	NO ₂	CH	-CH ₂ N(CH ₃)CH ₂ -	H	H	CH ₃
4.082	F	NO ₂	N	-CH ₂ N(CH ₃)CH ₂ -	H	H	CH ₃
4.083	Cl	CN	CH	-CH ₂ N(CH ₃)CH ₂ -	H	H	CH ₃
4.084	Cl	CN	N	-CH ₂ N(CH ₃)CH ₂ -	H	H	CH ₃
4.085	Cl	NO ₂	CH	-CH ₂ N(CH ₃)CH ₂ -	H	H	C ₂ H ₅
4.086	Cl	NO ₂	N	-CH ₂ N(CH ₃)CH ₂ -	H	H	C ₂ H ₅

4.087	F	NO ₂	CH	-CH ₂ N(CH ₃)CH ₂ -	H	H	C ₂ H ₅
4.088	Cl	NO ₂	CH	-CH ₂ N(CH ₃)CH ₂ -	CH ₃	H	C ₂ H ₅
4.089	Cl	CN	CH	-CH ₂ N(CH ₃)CH ₂ -	H	H	C ₂ H ₅
4.090	Cl	NO ₂	CH	-CH ₂ N(CH ₃)CH ₂ -	H	H	n-C ₃ H ₇
4.091	Cl	NO ₂	N	-CH ₂ N(CH ₃)CH ₂ -	H	H	n-C ₄ H ₉
4.092	F	NO ₂	CH	-CH ₂ N(CH ₃)CH ₂ -	H	H	s-C ₄ H ₉
4.093	Cl	NO ₂	CH	-CH ₂ N(CH ₃)CH ₂ -	H	H	t-C ₄ H ₉
4.094	Cl	CN	CH	-CH ₂ N(CH ₃)CH ₂ -	H	H	n-C ₇ H ₁₅
4.095	Cl	CN	N	-CH ₂ N(CH ₃)CH ₂ -	H	H	n-C ₁₁ H ₂₃
4.096	Cl	NO ₂	N	-(CH ₂) ₃ -	H	H	CH(CH ₃)C ₃ H ₇ -n
4.097	Cl	CN	N	-(CH ₂) ₃ -	H	H	C(CH ₃) ₂ C ₃ H ₇ -n
4.098	Cl	NO ₂	N	-(CH ₂) ₃ -	H	H	C(CH ₃) ₂ C ₂ H ₅
4.099	Cl	NO ₂	N	-(CH ₂) ₃ -	H	H	n-C ₆ H ₁₃
4.100	Cl	NO ₂	N	-(CH ₂) ₃ -	H	H	n-C ₁₀ H ₂₁
4.101	Cl	CN	C ₂ H ₅	CH ₃	H	H	CH(CH ₃)C ₃ H ₇ n
4.102	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	n-C ₅ H ₁₁
4.103	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	n-C ₈ H ₁₇
4.104	F	NO ₂	C ₂ H ₅	CH ₃	H	H	n-C ₄ H ₉
4.105	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	1-phenyl-1-cyclo- pentyl
4.106	Cl	CN	C ₂ H ₅	CH ₃	H	H	1-phenyl-1-cyclo- propyl
4.107	Cl	CN	C ₂ H ₅	CH ₃	H	H	1-phenyl-1-cyclo-

4.108	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	hexyl C(CH ₃) ₂ phenyl
4.109	Cl	CN	C ₂ H ₅	CH ₃	H	H	2,6-dimethylphenyl
4.110	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	2,4,6-tri-isopropyl- phenyl
4.111	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	cyclohexyl
4.112	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	benzyl
4.113	Cl	CN	C ₂ H ₅	CH ₃	H	H	1-adamantyl
4.114	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	t-C ₄ H ₉
4.115	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	CH(phenyl) ₂
4.116	F	NO ₂	C ₂ H ₅	CH ₃	H	H	cyclopropyl
4.117	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	CH(C ₂ H ₅) ₂
4.118	Cl	CN	C ₂ H ₅	CH ₃	H	H	CH(n-C ₃ H ₇) ₂
4.119	Cl	CN	C ₂ H ₅	CH ₃	H	H	CH ₂ (cyclohexyl)
4.120	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	CH ₂ CH ₂ (cyclo- hexyl)
4.121	Cl	CN	C ₂ H ₅	CH ₃	H	H	3-(t-C ₄ H ₉)cyclo- hex-1-yl
4.122	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	1-(4-chloro-phenyl)-1- cyclopentyl
4.123	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	1-(4-fluoro-phenyl)-1- cyclopentyl
4.124	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	CH ₂ C(CH ₃) ₂ CH ₃
4.125	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	CH ₂ CH(phenyl) ₂
4.126	Cl	CN	C ₂ H ₅	CH ₃	H	H	1-methyl-2,2-di- chloro-1-cyclopropyl
4.127	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	1-methyl-1-cyclohexyl
4.128	Cl	NO ₂	C ₂ H ₅	CH ₃	H	H	cyclopentyl
4.129	Cl	NO ₂	C ₂ H ₅	CH ₃	CH ₃	CH ₃	4-Cl-phenyl

Further typical and interesting compounds of formula (I) are:







Table 5: Compounds of formula (XXa)




No.	M	L	T	U	R ₂	R ₂ '	Hal	physical data
5.001	6-Cl	H	C ₂ H ₅	CH ₃	H	H	F	
5.002	6-Cl	5-Cl	C ₂ H ₅	CH ₃	H	H	F	
5.003	6-Cl	H	CH ₃	CH ₃	H	H	F	
5.004	6-Cl	H	C ₂ H ₅	CH ₃	CH ₃	H	F	
5.005	6-Cl	H	C ₂ H ₅	CH ₃	H	H	Cl	m.p. 142 - 143°C
5.006	6-Cl	H	C ₂ H ₅	CH ₃	H	H	Br	
5.007	6-Cl	H	C ₂ H ₅	CH ₃	H	H	I	

5.008	6-Cl	H	C ₂ H ₅	CH ₃	CH ₃	H	Cl
5.009	6-Cl	H	H	CH ₃	CH ₃	H	Br
5.010	H	5-Cl	C ₂ H ₅	CH ₃	H	H	Cl
5.011	6-Cl	H	C ₂ H ₅	CH ₃	H	H	Cl
5.012	6-Cl	H	C ₂ H ₅	H	H	H	Br
5.013	6-Cl	H	C ₂ H ₅	C ₂ H ₅	CH ₃	H	Br
5.014	6-Cl	H	C ₂ H ₅	CH ₃	C ₂ H ₅	H	F
5.015	6-Cl	H	C ₂ H ₅	CH ₃	n-C ₃ H ₇	H	F
5.016	6-Cl	H	C ₂ H ₅	CH ₃	s-C ₄ H ₉	H	F
5.017	6-Cl	H	C ₂ H ₅	CH ₃	n-C ₆ H ₁₃	H	F
5.018	6-Cl	H	CH ₃	CH ₃	CH ₃	H	F
5.019	6-Cl	4-F	CH ₃	CH ₃	CH ₃	H	F
5.020	6-Cl	H	C ₂ H ₅	CH ₃	CH ₃	CH ₃	I
5.021	6-Cl	H	C ₂ H ₅	CH ₃	CH ₃	CH ₃	F
5.022	6-Cl	H	C ₂ H ₅	CH ₃	C ₂ H ₅	CH ₃	Cl
5.023	6-Cl	H	C ₂ H ₅	CH ₃	CH ₃	CH ₃	Br
5.024	6-Cl	H	C ₂ H ₅	CH ₃	C ₂ H ₅	C ₂ H ₅	I

5.025	6-Cl	H	C ₂ H ₅	CH ₃	n-C ₃ H ₇	C ₂ H ₅	Cl
5.026	6-Cl	H	C ₂ H ₅	CH ₃	CH ₃	s-C ₄ H ₉	Br
5.027	6-Cl	H	C ₂ H ₅	CH ₃	CH ₃	n-C ₃ H ₇	I
5.028	6-Cl	H	C ₂ H ₅	CH ₃	s-C ₄ H ₉	C ₂ H ₅	F
5.029	6-Cl	H	C ₂ H ₅	CH ₃	s-C ₄ H ₉	n-C ₃ H ₇	Cl
5.030	6-Cl	H	C ₂ H ₅	C ₂ H ₅	CH ₃	n-C ₃ H ₇	I
5.031	6-Cl	H	C ₂ H ₅	CH ₃	C ₂ H ₅	CH ₃	F
5.032	6-Cl	H	C ₂ H ₅	CH ₃	n-C ₃ H ₇	CH ₃	F
5.033	6-Cl	H	C ₂ H ₅	CH ₃	s-C ₄ H ₉	CH ₃	F
5.034	6-Cl	H	C ₂ H ₅	CH ₃	n-C ₆ H ₁₃	H	F

Table 6: Compounds of formula (XXb)




No.	L	M	T	U	R ₂	R ₂ '	Hal
6.001	H	Cl	C ₂ H ₅	CH ₃	H	H	Cl
6.002	Cl	Cl	C ₂ H ₅	CH ₃	H	H	Cl

6.003	H	F	CH ₃	CH ₃	H	H	Cl
6.004	H	Cl	C ₂ H ₅	CH ₃	CH ₃	H	Cl
6.005	F	Cl	C ₂ H ₅	CH ₃	H	H	F
6.006	H	Cl	C ₂ H ₅	CH ₃	H	H	F
6.007	H	Cl	C ₂ H ₅	CH ₃	H	H	Br
6.008	H	Cl	C ₂ H ₅	CH ₃	CH ₃	H	F
6.009	H	Cl	H	CH ₃	H	H	Cl
6.010	H	Cl	C ₂ H ₅	C ₂ H ₅	H	H	Cl
6.011	H	Cl	C ₂ H ₅	CH ₃	H	H	I
6.012	H	Cl	C ₂ H ₅	CH ₃	CH ₃	H	I
6.013	H	Cl	C ₂ H ₅	CH ₃	CH ₃	CH ₃	I
6.014	H	H	C ₂ H ₅	C ₂ H ₅	CH ₃	H	F
6.015	H	Cl	C ₂ H ₅	CH ₃	C ₂ H ₅	H	Cl
6.016	H	Cl	C ₂ H ₅	CH ₃	n-C ₃ H ₇	H	Cl
6.017	H	Cl	C ₂ H ₅	CH ₃	s-C ₄ H ₉	H	Cl
6.018	H	Cl	C ₂ H ₅	CH ₃	n-C ₆ H ₁₃	H	Cl
6.019	H	Cl	CH ₃	CH ₃	CH ₃	H	Cl

6.020	H	F	CH ₃	CH ₃	CH ₃	H	Cl
6.021	H	Cl	C ₂ H ₅	CH ₃	C ₂ H ₅	CH ₃	I
6.022	H	Cl	C ₂ H ₅	CH ₃	CH ₃	H	Br
6.023	H	Cl	C ₂ H ₅	CH ₃	CH ₃	CH ₃	F

Table 7: Compounds of formula (XXc)




No.	Hal	Y	X	Q	R ₂	R ₂ '	Hal'
7.001	Cl	NO ₂	CH	H	H	H	Cl
7.002	Cl	NO ₂	N	H	H	H	Cl
7.003	F	NO ₂	CH	H	H	H	Cl
7.004	Cl	NO ₂	CH	3-CH ₃	H	H	Cl
7.005	Cl	CN	CH	H	H	H	Cl
7.006	Cl	CN	N	H	H	H	Cl
7.007	Cl	NO ₂	CH	H	H	H	F
7.008	Cl	NO ₂	N	H	H	H	F
7.009	F	NO ₂	CH	H	H	H	F

7.010	Cl	NO ₂	CH	2-CH ₃	H	H	F
7.011	Cl	CN	CH	H	H	H	Br
7.012	Cl	NO ₂	CH	H	H	H	Br
7.013	Cl	NO ₂	N	H	H	H	I
7.014	F	NO ₂	CH	H	H	H	I
7.015	Cl	NO ₂	CH	3-CH ₃	H	H	Cl
7.016	Cl	CN	CH	H	CH ₃	H	Cl
7.017	Cl	CN	N	H	CH ₃	H	Cl
7.018	Cl	NO ₂	CH	H	CH ₃	H	Cl
7.019	Cl	NO ₂	N	H	CH ₃	H	Cl
7.020	F	NO ₂	CH	H	CH ₃	H	Cl
7.021	Cl	NO ₂	CH	2-CH ₃	CH ₃	H	Cl
7.022	Cl	CN	CH	H	CH ₃	H	F
7.023	Cl	NO ₂	CH	H	CH ₃	H	F
7.024	Cl	NO ₂	N	H	CH ₃	CH ₃	F
7.025	F	NO ₂	CH	H	CH ₃	H	F

7.026	Cl	NO ₂	CH	3-CH ₃	CH ₃	H	F
7.027	Cl	CN	CH	H	CH ₃	H	Br
7.028	Cl	CN	N	H	CH ₃	H	Br
7.029	Cl	NO ₂	CH	3-C ₂ H ₅	H	H	Cl
7.030	Cl	CN	CH	2-CH ₃	CH ₃	CH ₃	Cl

Table 8: Compounds of formula (XXd)



No.	Hal	Y	X	T--U	R ₂	R ₂ '	Hal'
8.001	Cl	NO ₂	CH	-(CH ₂) ₃ -	H	H	Cl
8.002	Cl	NO ₂	N	-(CH ₂) ₃ -	H	H	Cl
8.003	F	NO ₂	CH	-(CH ₂) ₃ -	H	H	Cl
8.004	F	NO ₂	N	-(CH ₂) ₃ -	H	H	Cl
8.005	Cl	CN	CH	-(CH ₂) ₃ -	H	H	Cl
8.006	Cl	CN	N	-(CH ₂) ₃ -	H	H	Cl
8.007	Cl	NO ₂	CH	-(CH ₂) ₃ -	H	H	F
8.008	Cl	NO ₂	N	-(CH ₂) ₃ -	H	H	F
8.009	F	NO ₂	CH	-(CH ₂) ₃ -	H	H	F

8.010	Cl	NO ₂	CH	-(CH ₂) ₃ -	CH ₃	H	F
8.011	Cl	CN	CH	-(CH ₂) ₃ -	H	H	Br
8.012	Cl	NO ₂	CH	-(CH ₂) ₃ -	H	H	Br
8.013	Cl	NO ₂	N	-(CH ₂) ₃ -	H	H	I
8.014	F	NO ₂	CH	-(CH ₂) ₃ -	H	H	I
8.015	Cl	NO ₂	CH	-(CH ₂) ₃ -	CH ₃	H	Cl
8.016	Cl	CN	CH	-(CH ₂) ₃ -	CH ₃	H	Cl
8.017	Cl	CN	N	-(CH ₂) ₃ -	CH ₃	H	Cl
8.018	Cl	NO ₂	CH	-(CH ₂) ₃ -	CH ₃	H	Cl
8.019	Cl	NO ₂	N	-(CH ₂) ₃ -	CH ₃	CH ₃	Cl
8.020	F	NO ₂	CH	-(CH ₂) ₃ -	CH ₃	CH ₃	Cl
8.021	Cl	NO ₂	CH	-(CH ₂) ₃ -	CH ₃	CH ₃	Cl
8.022	Cl	CN	CH	-(CH ₂) ₃ -	CH ₃	H	F
8.023	Cl	NO ₂	CH	-(CH ₂) ₃ -	H	H	F
8.024	Cl	NO ₂	N	-(CH ₂) ₃ -	H	CH ₃	F
8.025	F	CN	CH	-(CH ₂) ₂ -	CH ₃	CH ₃	F

8.026	Cl	NO ₂	CH	-(CH ₂) ₂ -	CH ₃	CH ₃	F
8.027	Cl	CN	CH	-(CH ₂) ₃ -	H	H	Br
8.028	Cl	CN	N	-(CH ₂) ₃ -	H	H	Br
8.029	Cl	NO ₂	CH	-(CH ₂) ₃ -	C ₂ H ₅	H	Cl
8.030	Cl	CN	CH	-(CH ₂) ₃ -	C ₂ H ₅	H	Cl

It has now been found that systemic administration, for example by oral, percutaneous administration, or preferably topical application, for example in pour-on, spot-on or spray form, of a compound of formula (I) in an amount effective against insects is able drastically to reduce or completely to prevent attack by parasites on warm-blooded animals over a prolonged period.

The present invention therefore relates to the long-term control of parasites, especially blood-sucking insects but more especially the long-term control of fleas.

The compounds of formula (I) are distinguished *inter alia* by excellent activity against fleas, not only adult fleas being rapidly killed but also, by a circuitous route, the juvenile stages of the fleas. Flea larvae hatching out from the flea eggs feed mainly on the excreta of the adult fleas. Since the compounds of formula (I) according to the invention kill the adult fleas very rapidly, the necessary excreta are absent and the juvenile stages are deprived of the nutrient medium, so that they perish before reaching the adult stage.

The present invention therefore relates preferably to a method of controlling parasites on human beings, domestic animals, productive livestock and pets, which comprises administering systemically or preferably topically to the warm-blooded animal an effective amount of a composition comprising at least one compound of formula (I) or a physiologically tolerable salt thereof.

Long-term action is achieved by the compounds of formula (I) according to the invention using various forms of administration, for example by administering the active ingredient to

the animal to be treated externally or internally in a formulated form. "Formulated" in this case means, for example, in the form of a powder, a tablet or granules, in liposomes or a capsule, in the form of an emulsion, a foam or a spray, in microencapsulated form or in pour-on or spot-on form. It will be understood that all orally administrable compositions may comprise, in addition to customary formulation substances, further additives that encourage the host animal to take the composition orally voluntarily, e.g. suitable odorants and flavourings.

Percutaneous administration, for example by subcutaneous or intramuscular injection or as a depot preparation in the form of an implant, and topical application, for example in pour-on or spot-on form, represent preferred subjects of this invention on account of their being easy to carry out. A further mode of administration is oral administration, e.g. in the form of a tablet. Percutaneous and topical forms of administration are of particular interest and give excellent results.

Percutaneous forms of administration include, for example, subcutaneous, intramuscular and even intravenous administration of injectable forms. In addition to the customary syringes with needles, it is also possible to use needle-less high-pressure syringe devices.

Pour-on and spot-on formulations are especially preferred as topical forms of administration. but administration in the form of sprays, ointments, solutions or powders may also be expedient.

By selection of a suitable formulation it is possible to enhance the ability of the active ingredients to penetrate into the living tissue of the host animal and/or to maintain its availability. That is important when, for example, rather sparingly soluble active ingredients are used, the low solubility of which requires means for enhancing solubility, since in such cases the animal's body fluid will be capable of dissolving only small amounts of active ingredient at a time.

In order to obtain a greatly delayed release of active ingredient, a compound of formula (I) according to the invention may also be present in a matrix formulation which physically prevents the active ingredient from being released and excreted prematurely and maintains

the bioavailability of the active ingredient. Such a matrix formulation is injected into the body, e.g. intramuscularly or subcutaneously, and remains there as a form of depot from which active ingredient is released continuously. Such matrix formulations are known to a person skilled in the art. They are generally wax-like, semi-solid substances, for example vegetable waxes and polyethylene glycols having a high molecular weight, or solid polymer formulations, for example so-called microspheres.

The rate of release of the active ingredient from the implant and thus the period of time over which the implant exhibits an action is generally determined by the accuracy with which the implant has been calibrated (amount of active ingredient in the implant), the environment around the implant and the polymer formulation from which the implant has been made.

The administration of veterinary medicinal additives to animal food is well known in the field of animal health. It is usual first to prepare a so-called premix in which the active ingredient is dispersed in a liquid or is present in finely divided form in solid carriers. The premix can normally comprise about 1 to 800 mg of compound per kg of premix, depending on the desired final concentration in the food.

Since the compounds of formula (I) according to the invention may be hydrolysed by the constituents of the food, they should be formulated in a protective matrix, for example in gelatin, before being added to the premix.

The present invention accordingly relates also to the aspect of controlling parasites by administering to the host animal with its food a compound of formula (I) that has been protected against hydrolysis.

A compound of formula (I) according to the invention is advantageously administered in a dose of from 0.01 to 800 mg/kg, preferably from 0.1 to 200 mg/kg, especially from 0.5 to 30 mg/kg body weight, based on the host animal.

A good dose that can be administered regularly to the host animal is from 0.5 to 100 mg/kg, preferably from 0.1 to 40 mg/kg body weight. The administration is effected at suitable intervals in dependence upon the mode of administration and body weight.

The total dose may vary for the same active ingredient from one species of animal to another and also within a species of animal, since it depends *inter alia* on the weight, age and constitution of the host animal.

When used according to the invention, the compound of formula (I) according to the invention will normally be administered not in pure form but preferably in the form of a composition that comprises, in addition to the active ingredient, constituents that assist administration, suitable constituents being those which are tolerated by the host animal. It is of course possible, as well as controlling the adult parasites in accordance with the invention, additionally to use conventional methods to control the juvenile stages of the fleas, although the latter is not absolutely essential.

Such compositions to be administered in accordance with the invention generally comprise from 0.1 to 99 % by weight, especially from 0.1 to 95 % by weight, of a compound of formula (I) according to the invention and from 99.9 to 1 % by weight, especially from 99.9 to 5 % by weight, of a solid or liquid, physiologically tolerable carrier, including from 0 to 25 % by weight, especially from 0.1 to 25 % by weight, of a non-toxic dispersant.

Whereas commercial products will preferably be formulated as concentrates, the end user will normally employ dilute formulations.

Such formulations may also comprise further ingredients, such as stabilisers, antifoams, viscosity regulators, binders and tackifiers as well as other active ingredients for obtaining special effects.

The physiologically tolerable carriers known from veterinary medicinal practice for oral, percutaneous and topical administration can be used as formulation adjuvants. Some examples are given below.

Suitable carriers are especially fillers, such as sugars, for example lactose, saccharose, mannitol or sorbitol, cellulose preparations and/or calcium phosphates, for example tri-calcium phosphate or calcium hydrogen phosphate, and binders, such as starch pastes using, for example, maize, wheat, rice or potato starch, gelatin, tragacanth, methylcellulose

and/or, if desired, disintegrators, such as the above-mentioned starches, also carboxymethyl starch, cross-linked polyvinylpyrrolidone, agar, alginic acid or a salt thereof, such as sodium alginate. Adjuvants are especially flow conditioners and lubricants, for example silicic acid, talc, stearic acid or salts thereof, such as magnesium or calcium stearate, and/or polyethylene glycol. Dragée cores can be provided with suitable, optionally enteric, coatings, there being used *inter alia* concentrated sugar solutions which may comprise gum arabic, talc, polyvinylpyrrolidone, polyethylene glycol and/or titanium dioxide, or coating solutions in suitable organic solvents or solvent mixtures, or, for the preparation of enteric coatings, solutions of suitable cellulose preparations, such as acetylcellulose phthalate or hydroxypropylmethylcellulose phthalate. Dyes, flavourings or pigments may be added to the tablets or dragée coatings, for example for identification purposes or to indicate different doses of active ingredient.

Other orally administrable compositions are hard gelatin capsules, and also soft sealed capsules made of gelatin and a plasticiser, such as glycerol or sorbitol. The hard gelatin capsules may comprise the active ingredient in the form of granules, for example in admixture with fillers, such as lactose, binders, such as starches, and/or glidants, such as talc or magnesium stearate, and, optionally, stabilisers. In soft capsules, the active ingredient is preferably dissolved or suspended in suitable liquids, such as fatty oils, paraffin oil or liquid polyethylene glycols, to which stabilisers may also have been added. Preference is given *inter alia* to capsules that may easily be bitten through or swallowed without being chewed.

The pour-on or spot-on method comprises applying the compound of formula (I) to a locally limited area of the skin or coat, advantageously on the back of the neck or the backbone of the animal. This is carried out, for example, by applying the pour-on or spot-on formulation using a swab or spray to a relatively small area of the coat from where the active ingredient becomes distributed over a wide area of the coat almost automatically as a result of the spreading constituents of the formulation assisted by the movements of the animal.

Pour-on and spot-on formulations advantageously comprise carriers that assist rapid distribution over the surface of the skin and in the coat of the host animal and are generally

termed spreading oils. There are suitable, for example, oily solutions; alcoholic and isopropanolic solutions, e.g. solutions of 2-octyl dodecanol or oleyl alcohol; solutions in esters of monocarboxylic acids, such as isopropyl myristate, isopropyl palmitate, lauric acid oxalic ester, oleic acid oleyl ester, oleic acid decyl ester, hexyl laurate, oleyl oleate, decyl oleate, caproic acid esters of saturated fatty alcohols of chain length C_{12} - C_{18} ; solutions of esters of dicarboxylic acids, such as dibutyl phthalate, diisopropyl isophthalate, adipic acid diisopropyl ester, di-n-butyl adipate or solutions of esters of aliphatic acids, e.g. glycols. It may be advantageous for a dispersant known from the pharmaceutical or cosmetic industry also to be present. Examples are pyrrolidin-2-one, N-alkylpyrrolidin-2-one, acetone, polyethylene glycol and its ethers and esters, propylene glycol or synthetic triglycerides.

The oily solutions include e.g. vegetable oils, such as olive oil, groundnut oil, sesame oil, pine oil, linseed oil and castor oil. The vegetable oils may also be in epoxidised form. It is also possible to use paraffins and silicone oils.

Generally a pour-on or spot-on formulation will contain from 1 to 20 % by weight of a compound of formula (I), from 0.1 to 50 % by weight dispersant and from 45 to 98.9 % by weight solvent.

The pour-on and spot-on methods can be used especially advantageously for herd animals, such as cattle, horses, sheep and pigs, where it is difficult or time-consuming to treat all the animals orally or *via* injection. By virtue of its simplicity, this method can of course also be used for all other animals, including individual domestic animals and pets, and is welcomed especially by the keepers of the animals because it can frequently be carried out without the expert assistance of a veterinary surgeon.

Suitable for parenteral and percutaneous administration are oily injection solutions or suspensions, there being used suitable lipophilic solvents or vehicles, such as fatty oils, for example sesame oil, or synthetic fatty acid esters, for example ethyl oleate, or triglycerides, or aqueous injection solutions or suspensions that comprise viscosity-increasing substances, for example sodium carboxymethylcellulose, sorbitol and/or dextran, and, optionally, stabilisers.

The compositions of the present invention can be prepared in a manner known *per se*, for example by means of conventional mixing, granulating, confectioning, dissolving or lyophilising processes. For example, pharmaceutical compositions for oral administration can be obtained by combining the active ingredient with solid carriers, optionally granulating a resulting mixture, and processing the mixture or granules, if desired or necessary, after the addition of suitable excipients, to form tablets or dragée cores.

The following Examples and patent claims illustrate the invention described above, but do not limit its scope in any way. Temperatures are given in degrees Celsius. In the following Formulation Examples the expression "compound of formula (I)" is used to represent a compound of Tables 1 to 3, especially benzoic acid ({1-[(6-chloro-pyridin-3-ylmethyl)-ethyl-amino]-2-nitro-vinyl)-methyl-carbamoyloxy)-methyl ester.

Formulation Examples

Example 1: Tablets comprising 25 mg of a compound of formula (I) can be prepared as follows:

Constituents (for 1000 tablets)

compound of formula (I)	25.0 g
lactose	100.7 g
wheat starch	7.5 g
polyethylene glycol 6000	5.0 g
talcum	5.0 g
magnesium stearate	1.8 g
demineralised water	q.s.

Preparation: All the solid ingredients are first forced through a sieve of 0.6 mm mesh size. Then the active ingredient, the lactose, the talcum and half the starch are mixed together. The other half of the starch is suspended in 40 ml of water and the suspension is added to a boiling solution of the polyethylene glycol in 100 ml of water. The resulting starch paste is added to the main batch and the mixture is granulated, if necessary with the addition of water. The granules are dried overnight at 35°, forced through a sieve of 1.2 mm mesh size, mixed with the magnesium stearate and compressed to form tablets which have a mesh size of about 6 mm and which are concave on both sides.

Example 2: Injection solution

compound of formula (I)	0.1 to 10 %, preferably 0.5 to 5 %
non-ionic surfactant	0.1 to 30 %, preferably 0.5 to 10 %
mixture of ethanol and propylene glycol	60 to 99 %, preferably 85 to 90 %

Example 3: Injection suspension (aqueous or oily)

compound of formula (I)	0.1 to 20 %, preferably 1 to 10 %
non-ionic surfactant	0.1 to 20 %, preferably 1 to 10 %
water or vegetable oil	60 to 99 %, preferably 85 to 95 %

Example 4: Oily injectableA. Oily vehicle (slow release)

compound of formula (I)	0.1-1.0 g
groundnut oil	ad 100 ml

or

B.

compound of formula (I)	0.1-1.0 g
sesame oil	ad 100 ml

Preparation: The active ingredient is dissolved in a portion of the oil, with stirring and optionally with gentle heating, and after cooling the solution is made up to the desired volume and sterile-filtered through a suitable 0.22 mm membrane filter.

Example 5: Pour-on

A.

compound of formula (I)	10%
epoxidised soybean oil	5%
oleyl alcohol	85%

B.

compound of formula (I)	20%
pyrrolidin-2-one	15%
isopropyl myristate	65%

It is also possible to add to the described compositions biologically active substances or additives that have neutral behaviour towards the compounds of formula (I) and have no adverse effect on the host animal to be treated, and also mineral salts or vitamins.

Analogously to the described formulations of Examples 1 to 8 it is also possible to prepare further preparations having active ingredients of formula (I).

Example 6: Control of adult fleas in cats by means of pour-on application

Test protocol for each test compound: in order to determine the effectiveness of the test compounds against fully grown fleas, four groups each of two cats are used. Each cat is infested with 100 cat fleas [*Ctenocephalides felis* (Bouche)] and treated with 20 mg of active ingredient per kg body weight. The treatment is effected by applying the formulation to a locally limited area on the back of the cat's neck. While one group is infested with fleas but is treated only with a placebo, that is to say a formulation without active ingredient, and serves as control, another group is treated with nitenpyram as comparison substance; the two remaining groups receive the test compounds. Evaluation is made in each case by combing surviving fleas out of the animal's coat and comparing the number counted with the number of fleas in the control group and in the group treated with nitenpyram. The procedure in detail is as follows: each cat is infested with 100 fleas immediately after treatment on day 0. On day +1, each animal is combed and the number of surviving fleas is determined; the surviving fleas are then replaced on the same cat and after 24 hours the combing and evaluation are repeated. The fleas still surviving after those 24 hours are not returned to the cat. The described procedure is then repeated on days +3, +7, +9, +14, +21, +28, +35, +42, +49, +56 and +63 and in this way the effectiveness and duration of action are determined. The effectiveness is determined in accordance with the following formula:

$$\% \text{ effectiveness} = \frac{\text{number of living fleas on the control animal} - \text{number of living fleas on the test animal}}{\text{number of living fleas on the control animal}} \times 100$$

It is shown that the compounds of formula (I) according to the invention achieve excellent long-term action. Very good action is obtained, for example, with compounds Nos. 1.001, 1.008, 1.011, 1.012, 1.013, 1.014, 1.015, 1.018, 1.020, 1.021 and 1.022. An especially high level of long-term activity is exhibited by compounds 1.008, 1.011 and 1.012. Full action (100% effectiveness) is observed over a period of at least 7 weeks and after 7 weeks the action gradually declines to 26%. In comparison, nitenpyram exhibits full action (100% effectiveness) only for a maximum of 3 weeks. In a further, fully analogous test, compounds

1.019, 1.056, 1.057 and 1.058 also exhibit a long-term action equally as excellent as the compounds mentioned above.

In dogs the test proceeds in an entirely analogous manner. In order to investigate the total action and any side-effects after administration in the form of a pour-on formulation, compound No. 1.008 is selected and tested on 11 dogs. The test protocol and the results are given in Example 7.

Example 7: Control of adult fleas in dogs by means of pour-on application with compound No. 1.008 of Table 1

Test protocol for each test compound: in order to determine the effectiveness of the test compounds against fully grown fleas, three groups each of three dogs (1 male and 2 female beagles) are used; one group consists of two dogs (2 male beagles) and serves as control group. The entire test group consists of 11 beagles between 1 and 3.5 years of age. Each dog is infested with a total of 100 dog fleas [*Ctenocephalides canis*], 50 male and 50 female. Group 1 is treated with 2 mg, group 2 with 10 mg and group 3 with 20 mg of active ingredient per kg body weight. The treatment is effected by applying the pour-on formulation to a locally limited area on the back of the dog's neck. The fourth group is the control group. The latter is infested with fleas but is treated only with a placebo, that is to say a formulation without active ingredient, and serves for control purposes. Evaluation is made in each case by combing surviving fleas out of the animal's coat and comparing the number counted with the number of fleas in the control group. The procedure in detail is as follows: after treatment with the test formulation (day 0) each dog is infested on each of the subsequent days +1, +7, +14, +23, +28, +35, +49, +56, +63, +70, +77, +84 and +98 with 100 fleas. On the following day, each animal is combed and the number of surviving fleas is determined. The surviving fleas are then replaced on the same dog and after 24 hours the combing and evaluation are repeated. The fleas still surviving after those further 24 hours (2nd day after infestation with 100 fleas) are not returned to the dog. Then on days +3, +7, +9, +14, +21, +28, +35, +42 etc. the described procedure is repeated and in this way the effectiveness and the duration of action are determined. The effectiveness is determined using the formula given in Example 6.

79 days after application of the pour-on formulation, the test compound exhibits 100 % action; it is still 98.6 % effective on day 86 and even on day 98 it continues to be 92.3 % effective. Whereas the dogs from the three test groups exhibit no skin irritation or any other

undesirable side-effects, the two dogs in the control group have to be removed from the test programme on day 84, because they exhibit serious allergy symptoms and skin irritation as a result of the numerous flea bites.

Corresponding effects are also observed when the substances are administered not in pour-on form but in the form of an injection solution.

Example 8: Control of adult fleas in cats by means of subcutaneous injection


Test protocol for each test compound: in order to determine the effectiveness of the test compounds against fully grown fleas, four groups each of two cats from 1.5 to 4 years of age are used. Each cat is infested with 100 cat fleas [*Ctenocephalides felis* (Bouche)]. Two groups are treated with 20 mg of active ingredient per kg body weight. The treatment is effected by subcutaneous injection of a solution of the active ingredient behind the left shoulder blade. While one group is infested with fleas but is treated only with a placebo, that is to say a formulation without active ingredient, and serves as control, another group is treated with nitenpyram as comparison substance. Evaluation is made in each case by combing surviving fleas out of the animal's coat and comparing the number counted with the number of fleas in the control group and in the group treated with nitenpyram. The procedure in detail is as follows: each cat is infested with 100 fleas immediately after treatment on day 0. On day +1, each animal is combed and the number of surviving fleas is determined; the surviving fleas are then replaced on the same cat and after 24 hours the combing and evaluation are repeated. The fleas still surviving after those 24 hours are not returned to the cat. The described procedure is then repeated on days +3, +7, +9, +14, +21 and +28 and in this way the effectiveness and duration of action is determined. The effectiveness is determined using the same formula as in the preceding Example.

It is found that the compounds of formula (I) according to the invention achieve excellent long-term action after subcutaneous injection. Very good action is achieved, for example, with compounds Nos. 1.001, 1.008, 1.011, 1.012, 1.013, 1.014, 1.015, 1.018, 1.020, 1.021 and 1.022. An especially high level of long-term activity is exhibited by compound 1.012. Full action is observed for a period of at least 20 days in comparison with 2 days in the case of nitenpyram. The analogous test with dogs leads to absolutely comparable results.

Compounds 1.059 to 1.087 also exhibit very similar results to those described in Examples 6 to 8.

What is claimed is:

1. A compound of formula (I)



wherein

- R_1 is hydrogen or a radical from the group C_1 - C_4 alkyl, formyl, C_1 - C_6 alkylcarbonyl, C_1 - C_4 alkylsulfonyl, aryl, arylsulfonyl, arylcarbonyl, heterocyclyl and heterocyclyl-substituted C_1 - C_6 alkyl, which radical is unsubstituted or mono- or poly-substituted by identical or different substituents; the said substituents being C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 alkylthio, C_1 - C_4 haloalkyl, halogen, hydroxy, cyano, nitro, amino, C_1 - C_4 alkylamino, $(C_1$ - C_4 alkyl) $_2$ amino, alkoxy carbonyl, C_1 - C_4 alkylsulfonyl and arylsulfonyl;
- X is CH or N;
- Y is an electron-withdrawing radical;
- T has the meanings of R_1 or together with U forms a C_1 - C_4 alkylene bridge which is unsubstituted or substituted by a radical R_1 , or T and U together with the group -N-C-N- form a saturated or unsaturated 5- or 6-membered heterocyclic ring which may in addition contain as further hetero atom O or S or the hetero group -N(C_1 - C_6 alkyl)-;
- U is hydrogen or C_1 - C_6 alkyl;
- R_2 is hydrogen or C_1 - C_6 alkyl;
- R_2' is hydrogen or C_1 - C_6 alkyl; and
- R is C_1 - C_{20} alkyl, C_2 - C_{20} alkenyl, C_2 - C_6 alkynyl or heterocyclyl, each of those radicals being unsubstituted or substituted by one or more identical or different substituents, the said substituents being selected from the group halogen, cyano, nitro, hydroxy, C_1 - C_6 alkoxy, C_1 - C_6 alkylthio, C_1 - C_6 haloalkyl, C_1 - C_6 haloalkoxy and phenyl; or is C_3 - C_7 cycloalkyl that is unsubstituted or mono- or poly-substituted by identical or different substituents selected from halogen, cyano, nitro, hydroxy, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_1 - C_6 alkoxy, C_1 - C_6 alkylthio, C_1 - C_6 haloalkyl, C_1 - C_{20} haloalkoxy and phenyl;

wherein each phenyl moiety is itself unsubstituted or mono- or poly-substituted by identical or different substituents selected from halogen, cyano, nitro, hydroxy, C₁-C₆-alkyl, C₁-C₆alkoxy, C₁-C₆alkylthio, C₁-C₆haloalkyl and C₁-C₂₀haloalkoxy; or is phenyl phenoxyphenyl each of which is unsubstituted or mono- or poly-substituted by identical or different substituents selected from halogen, cyano, nitro, hydroxy, C₁-C₆alkyl, C₁-C₆alkoxy, C₁-C₆alkylthio, C₁-C₆haloalkyl, and C₁-C₂₀haloalkoxy.

2. A compound of formula (I) according to claim 1, wherein

Y is NO₂;

R₁ is hydrogen or a radical from the group C₁-C₄alkyl, formyl, C₁-C₆alkylcarbonyl, C₁-C₄-alkylsulfonyl, aryl, arylsulfonyl, arylcarbonyl, heterocyclyl and heterocyclyl-substituted C₁-C₆alkyl, which radical is unsubstituted or mono- or poly-substituted by identical or different substituents; the said substituents being C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄alkylthio, C₁-C₄haloalkyl, halogen, hydroxy, cyano, nitro, amino, C₁-C₄alkylamino, (C₁-C₄-alkyl)₂amino, alkoxycarbonyl, C₁-C₄alkylsulfonyl and arylsulfonyl;

T has the meanings of R₁ or together with U forms a C₁-C₄alkylene bridge which is unsubstituted or substituted by a radical R₁, or T and U together with the group -N-C-N- form a saturated or unsaturated 5- or 6-membered heterocyclic ring which may in addition contain as further hetero atom O or S or the hetero group -N(C₁-C₆alkyl)-;

U is hydrogen or C₁-C₆alkyl; and X, R₂, R₂' and R are as defined in claim 1 for formula (I).

3. A compound of formula (I) according to claim 1, wherein

R₁ is -CH₂-Het;

X is CH;

Y is NO₂;

Het is heterocyclyl that is unsubstituted or mono- or poly-substituted by identical or different substituents; the substituents being C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄alkylthio, C₁-C₄haloalkyl, halogen, hydroxy, cyano, nitro, amino, C₁-C₄alkylamino, (C₁-C₄alkyl)₂amino, alkoxycarbonyl, C₁-C₄alkylsulfonyl and arylsulfonyl;

T (1) is a radical from the group formyl, C₁-C₆alkylcarbonyl, C₁-C₄alkylsulfonyl, aryl, arylsulfonyl, arylcarbonyl, heterocyclyl and heterocyclyl-substituted C₁-C₆alkyl, which radical is unsubstituted or mono- or poly-substituted by identical or different substituents; the said substituents being C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkyl, halogen,


hydroxy, cyano, nitro, amino, C₁-C₄alkylamino, (C₁-C₄alkyl)₂amino, C₁-C₄alkylsulfonyl and arylsulfonyl; or

(2) T together with U forms a C₁-C₄alkylene bridge which is unsubstituted or substituted by a radical selected from the group C₁-C₄alkyl, formyl, C₁-C₆alkylcarbonyl, C₁-C₄alkylsulfonyl, aryl, arylsulfonyl, arylcarbonyl, heterocyclyl and heterocyclyl-substituted C₁-C₆alkyl; each radical from the said group itself being unsubstituted or substituted by C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkyl, halogen, hydroxy, cyano, nitro, amino, C₁-C₄alkylamino, (C₁-C₄alkyl)₂amino, C₁-C₄alkylsulfonyl or arylsulfonyl; or

(3) T and U together with the group -N-C-N- form a saturated or unsaturated 5- or 6-membered heterocyclic ring which may in addition contain as further hetero atom O or S or the hetero group -N(C₁-C₆alkyl)-;

U is hydrogen or C₁-C₆alkyl, preferably hydrogen, methyl or ethyl; and R₂, R₂' and R are as defined in claim 1 for formula (I).

4. A compound according to claim 1, which is a compound of formula (X)



wherein

R₁ is -CH₂-Het;

R is C₁-C₂₀alkyl, C₂-C₂₀alkenyl or C₂-C₆alkynyl, each of those radicals being unsubstituted or mono- or poly-substituted by identical or different substituents, the said substituents being selected from the group halogen, cyano, nitro, hydroxy, C₁-C₆alkoxy, C₁-C₆alkylthio, C₁-C₆haloalkyl, C₁-C₆haloalkoxy and phenyl; or is C₃-C₇cycloalkyl that is unsubstituted or mono- or poly-substituted by identical or different substituents selected from halogen, cyano, nitro, hydroxy, C₁-C₆alkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, C₁-C₆alkoxy, C₁-C₆alkylthio, C₁-C₆haloalkyl, C₁-C₂₀haloalkoxy and phenyl; or is phenyl phenoxy-phenyl each of which is unsubstituted or mono- or poly-substituted by identical or different substituents selected from halogen, cyano, nitro, hydroxy, C₁-C₆alkyl, C₁-C₆alkoxy, C₁-C₆alkylthio, C₁-C₆haloalkyl, C₁-C₂₀haloalkoxy;


T and U are each independently of the other hydrogen or C₁-C₆alkyl, preferably hydrogen, methyl or ethyl;

R₂ is hydrogen or C₁-C₆alkyl;

R_2' is hydrogen or C_1 - C_6 alkyl; and

Het is heterocyclyl that is unsubstituted or mono- or poly-substituted by identical or different halogen atoms.

5. A compound according to claim 1, which is a compound of formula (XI)



wherein


Hal is halogen, preferably fluorine, chlorine or bromine and especially chlorine; and especially occupies the 6-position in the pyridine;

X is CH or N;

Y is an electron-withdrawing radical;

T together with U forms a C_1 - C_4 alkylene bridge, which is unsubstituted or substituted by methyl or ethyl; and R_2 , R_2' and R are as defined in claim 1 for formula (I).

6. A compound according to claim 1, which is a compound of formula (XII)



wherein


Hal is halogen, preferably fluorine, chlorine or bromine and especially chlorine; and especially occupies the 2-position in the thiazole;

X is CH or N and especially N;


Y is an electron-withdrawing radical, preferably cyano, nitro or C_1 - C_6 haloalkyl-carbonyl, especially $CO-CF_3$; more especially nitro;

T together with U forms one of the groups $-CH_2-CH_2-$, $-CH_2-CH_2-CH_2-$, $-CH_2-O-CH_2-$ and $-CH_2-N(CH_3)-CH_2-$, wherein all methylene groups are unsubstituted or one of said methylene groups is substituted by methyl or ethyl; and R_2 , R_2' and R are as defined in claim 1 for formula (I).


7. A compound of formula (I) according to any one of claims 1 to 6, wherein U is methyl or ethyl; and the other substituents are as defined for formula (I) in claim 1.
8. A compound of formula (I) according to any one of claims 1 to 6, wherein T is methyl or ethyl; and the other substituents are as defined for formula (I) in claim 1.
9. A compound of formula (I) according to any one of claims 1 to 6, wherein R₂ and R₂' are each independently of the other hydrogen, methyl or ethyl; and the other substituents are as defined for formula (I) in claim 1.
10. A compound of formula (I) according to any one of claims 1 to 6, wherein heterocyclyl in R₁ is pyridyl, thiazolyl or tetrahydrofuryl that is unsubstituted or mono- or di-substituted by halogen; and the other substituents are as defined for formula (I) in claim 1.
11. A compound of formula (I) according to any one of claims 1 to 6, wherein heterocyclyl in R₁ is 5,6-dichloro-pyridin-3-yl, 6-chloro-pyridin-3-yl, 2-chlorothiazol-5-yl or tetrahydrofuran-3-yl, especially 6-chloro-pyridin-3-yl; and the other substituents are as defined for formula (I) in claim 1.
12. A process for the preparation of a compound of formula (I) according to any one of claims 1 to 11, which comprises either
- (a) reacting a compound of formula (IX)



in an aprotic, advantageously polar, solvent in the presence of a suitable base and at relatively low temperatures with a compound of formula (VIII)



- (b) reacting a compound of formula (XX)



with an acid RCOOH and isolating the end product from the reaction mixture, the substituents R, R₁, R₂, R₂', T, U, X and Y in formulae (IX) and (VIII) being as defined for formula (I), Hal being halogen, such as fluorine, chlorine, bromine or iodine, preferably chlorine, bromine or iodine, and W being a suitable leaving group.


13. A parasitocidal composition comprising a compound of formula (I) according to any one of claims 1 to 11 and at least one physiologically tolerable carrier.
14. A parasitocidal composition according to claim 13, comprising from 0.1 to 99 % by weight of a compound of formula (I) according to claim 1 and from 99.9 to 1 % by weight of a solid or liquid, physiologically tolerable carrier, including from 0 to 25 % by weight of a non-toxic dispersant.
15. A parasitocidal composition according to either claim 13 or claim 14, which is a pour-on or spot-on formulation.
16. A combination preparation for controlling parasites on warm-blooded animals, comprising, in addition to a compound of formula (I) according to any one of claims 1 to 11, at least one further active ingredient having the same or a different direction of action and at least one physiologically tolerable carrier.
17. A method of controlling parasites on warm-blooded animals, which comprises administering to a warm-blooded animal a parasitocidally effective compound of formula (I) according to any one of claims 1 to 11.
18. A method according to claim 17, comprising the topical application of a compound of formula (I) according to any one of claims 1 to 11.

19. A method according to claim 17, wherein a compound of formula (I) according to any one of claims 1 to 11 is administered in a dose of from 0.01 to 800 mg/kg body weight, based on the host animal.

20. The use of a compound of formula (I) according to any one of claims 1 to 11 in a method of controlling parasites of warm-blooded animals.

21. The use of a compound of formula (I) according to any one of claims 1 to 11 in the manufacture of a veterinary medicinal preparation against parasites.

22. A compound of formula (XX)



wherein

R₁ is hydrogen or a radical from the group C₁-C₄alkyl, formyl, C₁-C₆alkylcarbonyl, C₁-C₄-alkylsulfonyl, aryl, arylsulfonyl, arylcarbonyl, heterocyclyl and heterocyclyl-substituted C₁-C₆alkyl, which radical is unsubstituted or mono- or poly-substituted by identical or different substituents; the said substituents being C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄alkylthio, C₁-C₄haloalkyl, halogen, hydroxy, cyano, nitro, amino, C₁-C₄alkylamino, (C₁-C₄-alkyl)₂amino, alkoxycarbonyl, C₁-C₄alkylsulfonyl and arylsulfonyl;

X is CH or N;

Y is an electron-withdrawing radical, preferably cyano, nitro or C₁-C₆haloalkyl-carbonyl, especially CO-CF₃;

T has the meanings of R₁ or together with U forms a C₁-C₄alkylene bridge which is unsubstituted or substituted by a radical R₁, or T and U together with the group -N-C-N- form a saturated or unsaturated 5- or 6-membered heterocyclic ring which may in addition contain as further hetero atom O or S or the hetero group -N(C₁-C₆alkyl)-;


U is hydrogen or C₁-C₆alkyl, preferably hydrogen, methyl or ethyl;

R₂ is hydrogen or C₁-C₆alkyl;


R₂' is hydrogen or C₁-C₆alkyl; and

Hal is fluorine, chlorine, bromine or iodine, preferably chlorine.

23. A process for the preparation of a compound of formula (XX) according to claim 22, which comprises reacting a compound of formula (IX)



in an aprotic, advantageously polar, solvent in the presence of a suitable base and at relatively low temperatures with a compound of formula (VI)



the substituents R_1 , R_2 , R_2' , X, Y, T and U in formulae (XX), (IX) and (VI) being as defined for formula (I); W being a leaving group; and Hal being halogen, such as fluorine, chlorine, bromine or iodine, preferably chlorine, bromine or iodine.

24. The use of a compound of formula (XX) according to claim 22 in the preparation of a parasiticide.

25. A parasiticial composition comprising a compound of formula (XX) according to claim 22 and at least one physiologically tolerable carrier.

26. A method of controlling parasites on warm-blooded animals, which comprises administering to a warm-blooded animal a parasitically effective compound of formula (XX) according to claim 22.

27. The use of a compound of formula (XX) according to claim 22 in a method of controlling parasites on warm-blooded animals.

28. The use of a compound of formula (XX) according to claim 22 in the manufacture of a veterinary medicinal preparation against parasites.